

Spectral estimates for Schrödinger operator with periodic matrix potentials on the real line

Dmitri Chelkak ^{*}and Evgeny Korotyaev [†]

February 2, 2008

Abstract

We consider the Schrödinger operator on the real line with a $N \times N$ matrix valued periodic potential, $N > 1$. The spectrum of this operator is absolutely continuous and consists of intervals separated by gaps. We define the Lyapunov function, which is analytic on an associated N -sheeted Riemann surface. On each sheet the Lyapunov function has the standard properties of the Lyapunov function for the scalar case. The Lyapunov function has (real or complex) branch points, which we call resonances. We determine the asymptotics of the periodic, anti-periodic spectrum and of the resonances at high energy (in terms of the Fourier coefficients of the potential). We show that there exist two types of gaps: i) stable gaps, i.e., the endpoints are periodic and anti-periodic eigenvalues, ii) unstable (resonance) gaps, i.e., the endpoints are resonances (real branch points). Moreover, the following results are obtained: 1) we define the quasimomentum as an analytic function on the Riemann surface of the Lyapunov function; various properties and estimates of the quasimomentum are obtained, 2) we construct the conformal mapping with real part given by the integrated density of states and imaginary part given by the Lyapunov exponent. We obtain various properties of this conformal mapping, which are similar to the case $N=1$, 3) we determine various new trace formulae for potentials, the integrated density of states and the Lyapunov exponent, 4) a priori estimates of gap lengths in terms of potentials are obtained.

1 Introduction and main results

We consider the self-adjoint operator $\mathcal{L}y = -y'' + V(t)y$, acting in $L^2(\mathbb{R})^N$, $N \geq 2$ where the symmetric 1-periodic $N \times N$ matrix potential V belongs to the real Hilbert space $\mathcal{H}$ given by

$$\mathcal{H} = \left\{ V = V^* = \{V_{jk}(t)\}_{j,k=1}^N, \quad t \in \mathbb{R}/\mathbb{Z}, \quad \|V\|^2 = \int_0^1 \text{Tr } V^2(t) dt < \infty \right\}.$$

^{*}Dept. of Math. Analysis, Math. Mech. Faculty, St.Petersburg State University. Universitetskij pr. 28, Staryj Petergof, 198504 St.Petersburg, Russia, e-mail: delta4@math.spbu.ru

[†]Correspondence author. Institut für Mathematik, Humboldt Universität zu Berlin, Rudower Chaussee 25, 12489, Berlin, Germany, e-mail: evgeny@math.hu-berlin.de

It is well known (see [DS]) that the spectrum $\sigma(\mathcal{L})$ of $\mathcal{L}$ is absolutely continuous and consists of non-degenerated intervals $[\lambda_{n-1}^+, \lambda_n^-]$, $n = 1, \dots, N_G \leq \infty$. These intervals are separated by the gaps $\gamma_n = (\lambda_n^-, \lambda_n^+)$ with the length > 0 . **Without loss of generality we assume**

$$\lambda_0^+ = 0, \quad \text{and} \quad V^0 = \int_0^1 V(t)dt = \text{diag}\{V_1^0, V_2^0, \dots, V_N^0\}, \quad V_1^0 \leq V_2^0 \leq \dots \leq V_N^0. \quad (1.1)$$

A great number of papers is devoted to the inverse spectral theory for the Hill operator. We mention all papers where the inverse problem including characterization was solved: Marchenko and Ostrovski [MO1], Garnett and Trubowitz [GT1-2], Kappeler [Kap], Kargaev and Korotyaev [KK1], and Korotyaev [K1-3] and for 2×2 Dirac operator Misura [Mi1-2] and Korotyaev [K4-5]. Recently, the author [K6] extended the results of [MO1], [GT1], [K1-2] for the case $-y'' + uy$ to the case of distributions, i.e. $-y'' + u'y$ on $L^2(\mathbb{R})$, where periodic $u \in L^2_{loc}(\mathbb{R})$. It is important that in these papers new results from analytic function theory (in particular, conformal mapping theory) were obtained. As an example, we mention the proofs by the direct method (see [GT2], [KK1], [K1-3]). These are short, but this approach needs a priori estimates of potential in terms of spectral data. A priori estimates for various parameters of the Hill operator and for the Dirac operator (the norm of a periodic potential, effective masses, gap lengths, height of slits, and so on) were obtained in [GT1], [MO1-2], [KK1], [KK2], [K2-11]. In order to get the required estimates the authors of [GT1], [MO1-2], [Mi1-2], [K2-11]... used the “global quasi-momentum” (the conformal mapping), which was introduced into the spectral theory of the Hill operator by Marchenko-Ostrovski [MO1].

There exist many papers about the periodic systems $N \geq 2$ (see [Ca1-3], [YS]). The basic results for direct spectral theory for the matrix case were obtained by Lyapunov [Ly] (see also interesting papers of Krein [Kr], Gel'fand and Lidskii [GL]). In [BBK] for the case $N = 2$ the following results are obtained: the Lyapunov function is constructed on the 2-sheeted Riemann surface and the existence of real and complex branch points is proved. In [BK] the operator $y'''' + qy$, where q is a periodic real potential, was studied. In this case the Lyapunov function is constructed on a 2-sheeted Riemann surface and the existence of real and complex branch points is proved. The asymptotics of gaps and resonances in terms of the Fourier coefficients are obtained.

The main goal of our paper is to reformulate some spectral problem for the differential operator with periodic matrix coefficients as problems of conformal mapping theory. We construct the conformal mapping (averaged quasimomentum) w , with real part given by the integrated density of states and imaginary part given by the Lyapunov exponent. We obtain various properties of this conformal mapping, which are similar to the case $N = 1$. For solving these “new” problems we use some techniques from [KK2], [K2], [K6-8]. In particular, we use the Poisson integral for the domain $\mathbb{C}_+ \cup (-1, 1) \cup \mathbb{C}_-$ and the Dirichlet integral for the function $w(z) - z$. Note that the Dirichlet integral was used in [K6-8] for the scalar case to obtain a priori two-sided estimates of the potential in terms of spectral data.

Introduce the fundamental $N \times N$ -matrix solutions $\varphi(t, z)$, $\vartheta(t, z)$ of the equation

$$-f'' + V(t)f = z^2 f, \quad z \in \mathbb{C}, \quad (1.2)$$

with the conditions $\varphi(0, z) = \vartheta'(0, z) = 0$, $\varphi'(0, z) = \vartheta(0, z) = I_N$, where $I_N, N \geq 1$ is the identity $N \times N$ matrix. Here and below we use the notation $(') = \partial/\partial t$. We define the monodromy $2N \times 2N$ -matrix M and the trace $T_m, m \geq 1$ by

$$M(z) = \mathcal{M}(1, z), \quad \mathcal{M}(t, z) = \begin{pmatrix} \vartheta(t, z) & \varphi(t, z) \\ \vartheta'(t, z) & \varphi'(t, z) \end{pmatrix}, \quad T_m(z) = \frac{\text{Tr } M^m(z)}{2N}. \quad (1.3)$$

The functions $M(z)$ and $T_m, m \geq 1$ are entire, real for $z^2 \in \mathbb{R}$ and $\det M = 1$. Let $\tau_m, m = 1, \dots, 2N$ be the eigenvalues of M . An eigenvalue of $M(z)$ is called a *multiplier*. It is a root of the algebraic equation $D(\tau, z) \equiv \det(M(z) - \tau I_{2N}) = 0, \tau, z \in \mathbb{C}$. The zeros of $D(1, \sqrt{\lambda})$ (and $D(-1, \sqrt{\lambda})$ (counted with multiplicity) are the periodic (anti-periodic) eigenvalues for the equation $-y'' + Vy = \lambda y$ with periodic (anti-periodic) boundary conditions.

Below we need the following well-known results of Lyapunov (see [YS]).

Theorem (Lyapunov) *Let $V \in \mathcal{H}$. Then the following identities are fulfilled:*

$$M^{-1} = -JM^\top J = \begin{pmatrix} \varphi'(1, \cdot)^\top & -\varphi(1, \cdot)^\top \\ -\vartheta'(1, \cdot)^\top & \vartheta(1, \cdot)^\top \end{pmatrix}, \quad J = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix}, \quad (1.4)$$

$$D(\tau, \cdot) = \tau^{2N} D(\tau^{-1}, \cdot), \quad \tau \neq 0. \quad (1.5)$$

If for some $z \in \mathbb{C}$ (or $z^2 \in \mathbb{R}$) $\tau(z)$ is a multiplier of multiplicity $d \geq 1$, then $\tau^{-1}(z)$ (or $\bar{\tau}(z)$) is a multiplier of multiplicity d . Moreover, each $M(z), z \in \mathbb{C}$, has exactly $2N$ multipliers $\tau_m^{\pm 1}(z), m = 1, \dots, N$. Furthermore, $z^2 \in \sigma(V)$ iff $|\tau_m(z)| = 1$ for some $m = 1, \dots, N$. If $\tau(z)$ is a simple multiplier and $|\tau(z)| = 1$, then $\tau'(z) \neq 0$.

It is well known that $D(\tau, z) = \sum_0^{2N} \xi_m(z) \tau^{2N-m}$, where the functions ξ_m are given by

$$\xi_0 = 1, \quad \xi_1 = -2NT_1, \quad \xi_2 = -\frac{2N}{2}(T_2 + T_1\xi_1), \quad \dots, \xi_m = -\frac{2N}{m} \sum_0^{m-1} T_{m-j} \xi_j, \dots \quad (1.6)$$

see [RS]. Using the identity (1.5) we obtain

$$D(\tau, \cdot) = (\tau^{2N} + 1) + \xi_1(\tau^{2N-1} + \tau) + \dots + \xi_{N-1}(\tau^{N+1} + \tau^{N-1}) + \xi_N \tau^N. \quad (1.7)$$

The eigenvalues of $M(z)$ are the zeros of Eq. $D(\tau, z) = 0$. This is an algebraic equation in τ of degree $2N$. The coefficients $\xi_m(z)$ are entire in $z \in \mathbb{C}$. It is well known (see e.g. [Fo],[Sp]) that the roots $\tau_m(z), m = 1, \dots, 2N$ constitute one or several branches of one or several analytic functions that have only algebraic singularities in $\mathbb{C}$. Thus the number of eigenvalues of $M(z)$ is a constant N_e with the exception of some special values of z (see below the definition of a resonance). In general, there is a infinite number of such points on the plane. If the functions $\tau_m(z), m = 1, \dots, N$ are all distinct, then $N_e = 2N$. If some of them are identical, then we get $N_e < 2N$ and $M(z)$ is permanently degenerate.

The Riemann surface for the multipliers $\tau_m(z), m = 1, \dots, N$ has $2N$ sheets, since degree of $D(\tau, \cdot)$ is $2N$ see (1.7). If $N = 1$, then it has 2 sheets, but the Lyapunov function is entire. Similarly, in the case $N \geq 2$ it is more convenient for us to construct the Riemann surface

for the Lyapunov function, which has N sheets (see Eq. (1.9)). In order to formulate our first result we transform $D(\tau, z)$ to the polynomial $\Phi(\nu, z)$ by

$$\frac{D(\tau, z)}{(2\tau)^N} = \Phi(\nu, z) = \nu^N + \phi_1(z)\nu^{N-1} + \dots + \phi_N(z), \quad \nu = \frac{\tau + \tau^{-1}}{2}, \quad (1.8)$$

where $\phi_1, \dots, \phi_N$ are some linear combinations of $\xi_0, \dots, \xi_N$, see (2.12)-(2.15). In particular, all coefficients $\phi_1(z), \dots, \phi_N(z)$ are entire functions. Each zero of $\Phi(\nu, z)$ is a Lyapunov function

$$\Delta_m(z) = \frac{1}{2}(\tau_m(z) + \tau_m^{-1}(z)), \quad m = 1, \dots, N.$$

Remark. We note that this reduction from the polynomial D with $\deg D = 2N$ to the polynomial Φ with $\deg \Phi = N$ is crucial for our analysis. It is based on (1.5), which is a consequence of M being a symplectic matrix and on the identity (2.11) for the Chebyshev polynomials.

We need the following preliminary results

Theorem 1.1. *Let $V \in \mathcal{H}$. Then there exist analytic functions $\tilde{\Delta}_s, s = 1, \dots, N_0 \leq N$ on the N_s -sheeted Riemann surface $\mathcal{R}_s, N_s \geq 1$ having the following properties:*

i) There exist disjoint subsets $\omega_s \subset \{1, \dots, N\}, s = 1, \dots, N_0, \bigcup \omega_s = \{1, \dots, N\}$ such that all branches of $\tilde{\Delta}_s, s = 1, 2, \dots, N_0$ have the form $\Delta_j(z) = \frac{1}{2}(\tau_j(z) + \tau_j^{-1}(z)), j \in \omega_s$. Moreover, for any $z, \tau \in \mathbb{C}$ the following identities are fulfilled:

$$\frac{D(\tau, z)}{(2\tau)^N} = \prod_1^{N_0} \Phi_s(\nu, z), \quad \Phi_s(\nu, z) = \prod_{j \in \omega_s} (\nu - \Delta_j(z)), \quad \nu = \frac{\tau + \tau^{-1}}{2}, \quad \tau \neq 0, \quad (1.9)$$

where the functions $\Phi_s(\nu, z)$ are entire with respect to $\nu, z \in \mathbb{C}$ and $\Phi_s(\nu, z) \in \mathbb{R}$ for all $\nu, z \in \mathbb{R}$. Moreover, if $\Delta_i = \Delta_j$ for some $i \in \omega_k, j \in \omega_s$, then $\Phi_k = \Phi_s$ and $\tilde{\Delta}_k = \tilde{\Delta}_s$.

ii) (The monotonicity property). Let some branch $\Delta_m, m = 1, \dots, N$ be real analytic on some interval $Y = (\alpha, \beta) \subset \mathbb{R}$ and $-1 < \Delta_m(z) < 1$ for any $z \in Y$. Then $\Delta'_m(z) \neq 0$ for each $z \in Y$.

iii) Each function $\rho_s, s = 1, \dots, N_0$ given by (1.10) is entire and real on the real line,

$$\rho = \prod_1^{N_0} \rho_s, \quad \rho_s(\cdot) = \prod_{i < j, i, j \in \omega_s} (\Delta_i(\cdot) - \Delta_j(\cdot))^2. \quad (1.10)$$

iv) Each gap $\gamma_n = (\lambda_n^-, \lambda_n^+), n \geq 1$ is a bounded interval and $\lambda_n^\pm$ are either periodic (anti-periodic) eigenvalues or real branch points of Δ_m (for some $m = 1, \dots, N$) which are zeros of ρ (below we call such points resonances).

Remark. 1) If $N_0 = N$, then $\mathcal{R}_m = \mathbb{C}$ and each function $\Delta_m, m = 1, \dots, N$ is entire.

2) We have the following asymptotics (see Sect. 3)

$$\Delta_m(z) = \cos z + \frac{\sin z}{2z} V_m^0 + O\left(\frac{e^{|\operatorname{Im} z|}}{z^2}\right), \quad m = 1, \dots, N, \quad |z| \rightarrow \infty. \quad (1.11)$$

Then firstly, ρ is not a polynomial since ρ is bounded on $\mathbb{R}$. Secondly, if $V_i^0 \neq V_j^0, i \neq j$, then (1.11) implies $\Delta_i \neq \Delta_j$.

3) Let the surface $\mathcal{R} = \cup_1^{N_0} \mathcal{R}_s$ be a union of the disjoint Riemann surfaces $\mathcal{R}_s$ and let $\Delta = \{\tilde{\Delta}_s, s = 1, \dots, N_0\}$ be the corresponding analytic function on $\mathcal{R}$. Let $\phi : \mathcal{R} \rightarrow \mathbb{C}$ be the projection from the surface $\mathcal{R}$ into the complex plane. **We set** $\zeta \in \mathcal{R}$ and $z \in \mathbb{C}$.

Definition. The number z_0 is a **resonance** of $\mathcal{L}$, if z_0 is a zero of ρ given by (1.10).

Define the real matrices $V^{sn} = \{V_{jk}^{sn}\}, V^{cn} = \{V_{jk}^{cn}\}$ by

$$\hat{V}^{(n)} = \{\hat{V}_{jk}^{(n)}\} = \hat{V}^{cn} + i\hat{V}^{sn} = \int_0^1 V(t) e^{i2\pi nt} dt. \quad (1.12)$$

Denote by $\lambda_m^{n,\pm}, n \geq 0, m \in \{1, 2, \dots, N\}$ the eigenvalues of the periodic and anti-periodic problem for the equation $-f'' + Vf = z^2 f$. The periodic eigenvalues (n is even) satisfy

$$0 \leq \underbrace{\lambda_1^{0,+} \leq \lambda_2^{0,+} \leq \dots \leq \lambda_N^{0,+}}_{n=0} \leq \underbrace{\lambda_1^{2,-} \leq \lambda_1^{2,+} \leq \dots \leq \lambda_N^{2,-} \leq \lambda_N^{2,+}}_{n=2} \leq \lambda_1^{4,-} \leq \lambda_1^{4,+} \leq \dots \quad (1.13)$$

Recall $\lambda_0^+ = 0$. The anti-periodic eigenvalues (n is odd) satisfy

$$0 \leq \underbrace{\lambda_1^{1,-} \leq \lambda_1^{1,+} \leq \dots \leq \lambda_N^{1,-} \leq \lambda_N^{1,+}}_{n=1} \leq \underbrace{\lambda_1^{3,-} \leq \lambda_1^{3,+} \leq \dots \leq \lambda_N^{3,-} \leq \lambda_N^{3,+}}_{n=3} \leq \lambda_1^{5,-} \leq \dots \quad (1.14)$$

If $V = 0$, then $\lambda_m^{n,\pm} = (\pi n)^2, m = 1, \dots, N$. Let $z_m^{n,\pm} = \sqrt{\lambda_m^{n,\pm}} > 0$ and $z_m^{-n,\pm} = -z_m^{n,\mp}, n \geq 0, m \in \{1, 2, \dots, N\}$. The zeros of $D(1, z)$ and $D(-1, z)$ (counted with multiplicity) have the forms $z_m^{2n,\pm}$ and $z_m^{2n+1,\pm}, n \in \mathbb{Z}$. Let $|A|$ denote the operator norm of the matrix A .

Theorem 1.2. Let $V \in \mathcal{H}$. Then the periodic and anti-periodic eigenvalues have the following asymptotics:

$$\lambda_m^{n,\pm} = (\pi n)^2 + \zeta_m^{n,\pm} + O(n^{-1}), \quad m = 1, \dots, N, \quad n \rightarrow \infty, \quad (1.15)$$

where $\zeta_m^{n,\pm}, m = 1, 2, \dots, N$ are the eigenvalues of the matrix $\begin{pmatrix} V^0 + \hat{V}^{cn} & \hat{V}^{sn} \\ \hat{V}^{sn} & V^0 - \hat{V}^{cn} \end{pmatrix}$.

Assume that $V_j^0 \neq V_{j'}^0$ for all $j \neq j' \in \omega_s$ for some $s = 1, \dots, N_0$. Then the function ρ_s has the zeros $z_\alpha^{n,\pm}, \alpha = (j, j'), j < j', j, j' \in \omega_s, n \in \mathbb{Z} \setminus \{0\}$, which are real at large n and satisfy

$$z_\alpha^{n,\pm} = \pi n + \frac{V_i^0 + V_j^0}{4\pi n} + O\left(\frac{|\hat{V}^{(n)}|}{n} + \frac{1}{n^2}\right), \quad \alpha = (j, j') \quad \text{as } n \rightarrow \infty. \quad (1.16)$$

Let in addition $V_1^0 < \dots < V_N^0$. Then for each $s = 1, \dots, N_0$ and for large $n \rightarrow \infty$ there exists a system of real intervals (gaps) $\gamma_\alpha^n = (\lambda_\alpha^{n-}, \lambda_\alpha^{n+}), g_\alpha^n = (z_\alpha^{n-}, z_\alpha^{n+})$, such that

$$z_{j,j'}^{n,\pm} = z_{j',j}^{n,\pm} > 0, \quad \lambda_{j,j}^{n,\pm} = \lambda_j^{n,\pm}, \quad (z_\alpha^{n,\pm})^2 = \lambda_\alpha^{n,\pm} > 0, \quad \alpha = (j, j'), \quad j, j' \in \omega_s,$$

$$\lambda_{j,j_1}^{n-} \leq \lambda_{j,j_1}^{n+} < \lambda_{j,j_2}^{n-} \leq \lambda_{j,j_2}^{n+} < \dots < \lambda_{j,j_{N_s}}^{n-} \leq \lambda_{j,j_{N_s}}^{n+}, \quad N_s = |\omega_s|$$

$$(-1)^n \Delta_j(z) > 1, z \in g_{j,j}^n, \quad \text{and} \quad \overline{\Delta}_{j'}(z) = \Delta_j(z), z \in g_{j,j'}^n, \quad \text{if } j \neq j' \quad (1.17)$$

i) The branch Δ_j is real and is analytic on the set $(\pi n - \frac{\pi}{2}, \pi n + \frac{\pi}{2}) \setminus \cup_{p \neq j} g_{j,p}^n$ and is not real on $\cup_{p \neq j} g_{j,p}^n$.

ii) If $z_\alpha^{n-} \neq z_\alpha^{n+}$ for some $\alpha = j, j', j \neq j'$, then $z_\alpha^{n\pm}$ is a simple branch point, i.e., of square root type (resonance) for the functions $\Delta_j, \Delta_{j'}$. If $z_\alpha^{n-} = z_\alpha^{n+}$, then $\Delta_j, \Delta_{j'}$ are analytic at $z_\alpha^{n\pm}$.

iii) The following asymptotics are fulfilled:

$$\lambda_\alpha^{n\pm} = (\pi n)^2 + \frac{V_i^0 + V_j^0}{2} \pm |\hat{V}_\alpha^{(n)}| + O\left(|\hat{V}^{(n)}| + \frac{1}{n}\right), \quad \alpha = (j, j'). \quad (1.18)$$

Remark. 1) If $N = 1$, then the asymptotics (1.18) are well known [Ti]. 2) We describe the surface $\mathcal{R}$ in the case $V_1^0 < \dots < V_N^0$ and $N_0 = 1$ for large z . To "build" the surface $\mathcal{R}$ for large z , we take N replicas of the z -plane and call them sheets $\mathcal{R}_1, \dots, \mathcal{R}_N$. Each $\mathcal{R}_j$ is cut along the real interval $g_\alpha^n, \alpha = (j, j'), j, j' = 1, \dots, N, j \neq j'$. The cut on each sheet has two edges; we label each edge with $\sharp$ or $\flat$. Then attach the $\sharp$ edge of the cut g_α^n on $\mathcal{R}_j$ to the $\flat$ edge of the same cut on $\mathcal{R}_{j'}, j \neq j'$, and attach the $\flat$ edge of the cut on $\mathcal{R}_j$ to the $\sharp$ edge of the same cut on $\mathcal{R}_{j'}$. Thus, whenever we cross the cut, we pass from one sheet to the other. 3) If $V_1^0 < \dots < V_N^0$, then all resonances are real at high energy. The existence of low energy complex resonances for specific potentials was established in [BKK].

Recall that N_G is the total number of gaps in the spectrum of $\mathcal{L}$.

Corollary 1.3. Let $V \in \mathcal{H}$ and $V_1^0 < \dots < V_N^0$.

(i) If the identity $V_1^0 + V_N^0 = V_2^0 + V_{N-1}^0 = \dots = V_N^0 + V_1^0$ is not fulfilled, then $N_G < \infty$.

(ii) If $V_1^0 + V_N^0 = \dots = V_N^0 + V_1^0$ holds true and there exists a sequence of indices $n_k \rightarrow \infty$ such that $|\hat{V}^{(n_k)}|^2 + n_k^{-1} = o(|\hat{V}_{m, N+1-m}^{(n_k)}|)$, $k \rightarrow \infty$, for each $m = 1, \dots, N$, then $N_G = \infty$.

Remark. 1) Maksudov and Veliev [MV] proved i), for $N \geq 3$ in a more general case.

2) If $V = \text{diag}\{V_{11}, V_{22}, \dots, V_{NN}\}$ and $V_1^0 < V_2^0$, then the number of gaps is $N_G < +\infty$.

3) Note that the condition $|\hat{V}^{(n)}|^2 + n^{-1} = o(|\hat{V}_{m, N+1-m}^{(n)}|)$, $m = 1, \dots, N$, $n \rightarrow \infty$, holds true for "generic" potentials from the space $\mathcal{H}$. This yields the existence of real resonance gaps $(\lambda_\alpha^{n-}, \lambda_\alpha^{n+})$ at high energy. The coefficients $V_{m, N+1-m}, m = 1, \dots, N$ (the "second diagonal" of the matrix V) "create" the gaps.


We consider the conformal mapping associated with the operator $\mathcal{L}$. We need functions from the subharmonic counterpart of the Cartwright class of the entire functions given by

$$\mathcal{SC} = \left\{ v : \mathbb{C} \rightarrow \mathbb{R}, \quad \begin{array}{l} v \text{ is subharmonic in } \mathbb{C} \text{ and harmonic outside } \mathbb{R}, \\ v(\bar{z}) \equiv v(z), \int_{\mathbb{R}} \frac{v_+(t) dt}{1+t^2} < \infty, \limsup_{z \rightarrow \infty} \frac{v(z)}{|z|} < \infty \end{array} \right\}. \quad (1.19)$$


We recall the class of functions from [KK1] given by

$$\mathcal{SK}_m^+ = \left\{ v \in \mathcal{SC} : v \geq 0, \quad \lim_{y \rightarrow \infty} \frac{v(iy)}{y} = 1, \quad \int_{\mathbb{R}} (1+t^{2m})v(t)dt < \infty \right\}, \quad m \geq 0.$$

Model of 3-sheeted Riemann surface $\mathcal{R}$ near points πn and $\pi(n+1)$:



Real branches of 3-valued Lyapunov function $\Delta(x)$:



Gaps in spectrum:




Image of averaged quasi-momentum, w -plane:




Figure 1: $N = 3$, $V_1^0 < V_2^0 < V_3^0$, $\frac{1}{2}(V_1^0 + V_3^0) < V_2^0$.

We note that $\mathcal{SK}_{m+1}^+ \subset \mathcal{SK}_m^+, m \geq 0$.

Introduce the simple conformal mapping $\eta : \mathbb{C} \setminus [-1, 1] \rightarrow \{\zeta \in \mathbb{C} : |\zeta| > 1\}$ by

$$\eta(z) = z + \sqrt{z^2 - 1}, \quad z \in \mathbb{C} \setminus [-1, 1], \quad \text{and} \quad \eta(z) = 2z + o(1), \quad |z| \rightarrow \infty. \quad (1.20)$$

Note that $\eta(z) = \bar{\eta}(\bar{z}), z \in \mathbb{C} \setminus [-1, 1]$ since $\eta(z) > 1$ for any $z > 1$. Due to the properties of the Lyapunov functions we have $|\eta(\tilde{\Delta}_s(\zeta))| > 1, \zeta \in \mathcal{R}_s^+ = \{\zeta \in \mathcal{R}_s : \text{Im } \zeta > 0\}$. Thus we can introduce the quasimomentum k_m (we fix some branch of arccos and $\Delta_m(z)$) and the function q_m by

$$k_m(z) = \arccos \Delta_m(z) = i \log \eta(\Delta_m(z)), \quad q_m(z) = \text{Im } k_m(z) = \log |\eta(\Delta_m(z))|, \quad (1.21)$$

$m = 1, 2, \dots, N$ and $z \in \mathcal{R}_0^+ = \mathbb{C}_+ \setminus \beta_+, \beta_+ = \bigcup_{\beta \in \mathcal{B}_\Delta \cap \mathbb{C}_+} [\beta, \beta + i\infty)$ where $\mathcal{B}_\Delta$ is the set of all branch points of the function Δ . The branch points of k_m belong to $\mathcal{B}_\Delta$. Define the **averaged quasimomentum** w , the **density** u and the **Lyapunov exponent** v by

$$w(z) = u(z) + iv(z) = \frac{1}{N} \sum_1^N k_m(z), \quad v(z) = \text{Im } w(z), \quad z \in \mathcal{R}_0^+. \quad (1.22)$$

Define the sets $\sigma_{(N)} = \{z \in \mathbb{R} : \Delta_1(z), \dots, \Delta_N(z) \in [-1, 1]\}$ and

$$\sigma_{(1)} = \{z \in \mathbb{R} : \Delta_m(z) \in (-1, 1), \Delta_p(z) \notin [-1, 1] \text{ some } m, p = 1, \dots, N\}.$$

For the function $w(z) = u(z) + iv(z), z = x + iy \in \overline{\mathbb{C}}_+$ we formally introduce the integrals

$$Q_n = \frac{1}{\pi} \int_{\mathbb{R}} x^n v(x) dx, \quad \mathcal{P}_n = \frac{1}{\pi} \int_{\mathbb{R}} x^n v(x) du(x), \quad I_n^D = \frac{1}{\pi} \iint_{\mathbb{C}_+} |w'_n(z)|^2 dx dy, \quad (1.23)$$

$n \geq 0$, where here and below $w_m(z), z \in \mathbb{C}_+$ is given by

$$w_m(z) = \frac{1}{\pi} \int_{\mathbb{R}} \frac{t^m v(t)}{t - z} dt = z^m \left(w(z) - z + \sum_{n=0}^{m-1} Q_n z^{-n-1} \right), \quad z \in \mathbb{C}_+.$$

Let C_{us} denote the class of all real upper semi-continuous functions $h : \mathbb{R} \rightarrow \mathbb{R}$. With any $h \in C_{us}$ we associate the "upper" domain $W(h) = \{w = u + iv \in \mathbb{C} : v > h(u), u \in \mathbb{R}\}$. Let $g = \bigcup_{n \in \mathbb{Z}} g_n$ where $g_n = (z_n^-, z_n^+), z_n^\pm = \sqrt{\lambda_n^\pm} > 0$ and $g_{-n} = g_n, n \geq 1$. We formulate our main result.

Theorem 1.4. *Let $V \in \mathcal{H}$. Then the averaged quasimomentum $w = \frac{1}{N} \sum_1^N k_m$ is analytic in $\mathbb{C}_+$ and $w : \mathbb{C}_+ \rightarrow w(\mathbb{C}_+) = W(h)$ is a conformal mapping onto $W(h)$ for some $h \in C_{us}$. Moreover, $v = \text{Im } w$ has an harmonic extension from $\mathbb{C}_+$ into $\Omega = \mathbb{C}_+ \cup \mathbb{C}_- \cup g$ given by $v(z) = v(\bar{z}), z \in \mathbb{C}_-$ and $v(z) > 0$ for any $z \in \Omega$. Furthermore $v \in \mathcal{SK}_2^+ \cap C(\mathbb{C})$ and there exist branches $k_m, m = 1, \dots, N$ such that the following asymptotics, identities and estimates are fulfilled:*

$$w(z) = -\overline{w(-\bar{z})}, \quad z \in \overline{\mathbb{C}}_+, \quad (1.24)$$

$$w(z) - z = -\frac{Q_0}{z} - \frac{Q_2 + o(1)}{z^3}, \quad \text{as } y > r|x|, \quad y \rightarrow \infty, \quad \text{for any } r > 0, \quad (1.25)$$

$$Q_0 = I_0^D + \mathcal{P}_0 = \int_0^1 \frac{\text{Tr } V(t) dt}{2N}, \quad (1.26)$$

$$Q_2 = I_1^D + \mathcal{P}_2 = \int_0^1 \frac{\text{Tr } V^2(t) dt}{2^3 N}, \quad (1.27)$$

$$v|_{\sigma(N)} = 0, \quad 0 < v|_{\sigma(1) \cup g} \leq \sqrt{2Q_0}, \quad (1.28)$$

Remark. 1) Craig and Simon [CS] proved that the Lyapunov exponent is subharmonic in $\mathbb{C}$ for the Schrödinger operator $-\frac{d^2}{dx^2} + V$ for a large class of scalar potentials.
2) Similar arguments give

$$Q_4 = I_2^D + \mathcal{P}_4 = \int_0^1 \frac{\text{Tr}(V'(t)^2 + 2V^3(t)) dt}{2^5 N}, \quad \text{if } V' \in \mathcal{H}. \quad (1.29)$$

3) The integral $\mathcal{P}_0 \geq 0$ is the area between the boundary of $W(h)$ and the real line. The mapping $w : \mathbb{C}_+ \rightarrow W(h)$ is illustrated in Figure 2. In Figure 1 the upper picture is a domain $W(h)$ and $\tilde{A} = w(A), \tilde{B} = w(B), \dots$. The spectral interval (A, B) (with multiplicity 2) of the z -domain is mapped on the curve $(\tilde{A}, \tilde{B})$ of the w -domain, the interval (a gap) (B, C) of the z -domain is mapped on a vertical slit, which lies on the line $\text{Re } w = \pi$. The spectral interval (C, D) (with multiplicity 2) of the z -domain is mapped on the curve $(\tilde{C}, \tilde{D})$ of the w -domain. The spectral interval (D, E) (with multiplicity 4) of the z -domain is mapped on the interval $(\tilde{D}, \tilde{C})$ of the w -domain. The case of the interval (E, J) is similar. The resonance gap (K, L) of the z -domain is mapped on the vertical slit on the line $\text{Re } w = 3\pi$. In fact the boundary of $W(h)$ is given by the graph of the function $h(u), u \in \mathbb{R}$.




Figure 2: $N = 2$. The domain $W(h) = w(\mathbb{C}_+)$ and gaps in the spectrum.

Let $\sigma(m, V)$ denote the spectrum of $\mathcal{L}$ of multiplicity $2m, m \geq 0$. We have the following simple corollary from Theorem 1.4.

Corollary 1.5. *Let $\sigma(\mathcal{L}) = \sigma(N, V) = \mathbb{R}_+$ for some $V \in \mathcal{H}$. Then $V = 0$.*

Proof. Due to $\sigma_{(N)} = \mathbb{R}_+$, we obtain $Q_0 = Q_2 = 0$. Then identity (1.26) yields $\|V\| = 0$. ■

Recall that in the scalar case, the so-called Borg Theorem follows immediately from the existence of the conformal mapping and the asymptotics of the Lyapunov function at high energy [M]. In general, in order to prove the uniqueness result (the simplest part in the inverse spectral theory) it is necessary to use some results from "function theory". In our case we use conformal mapping theory. Recall that the so-called Borg Theorem for periodic systems was proved in [CHGL],[GKM] for general cases.

We describe the properties of the conformal mapping w .

Theorem 1.6. *Let $V \in \mathcal{H}$. Then the following relations are fulfilled:*

$$u'_x(z) \geq 1, \quad z \in \sigma_{(N)} \quad \text{and} \quad u'_x(z) > 0, \quad z \in \sigma_{(1)}, \quad (1.30)$$

here $u'_x(z) = 1$ for some $z \in \sigma_{(N)}$ iff $V = 0$. Moreover,

$$v''_{xx}(z) < 0 < v(z), \quad u(z) = \text{const} \in \frac{\pi}{N}\mathbb{Z}, \quad \text{for all } z \in g_n = (z_n^-, z_n^+), \quad (1.31)$$

$$v(x) = v_n^0(x) \left(1 + \frac{1}{\pi} \int_{\mathbb{R} \setminus g_n} \frac{v(t)dt}{v_n^0(t)|t-x|} \right), \quad x \in g_n, \quad v_n^0(z) = |(z - z_n^-)(z_n^+ - z)|^{\frac{1}{2}}, \quad (1.32)$$

$$\sum_n |g_n|^2 \leq 8Q_0, \quad G^2 \equiv \sum |\gamma_n|^2 \leq \frac{8}{N} \|V\|^2, \quad (1.33)$$

$$\|V\| \leq C_0 G (1 + G^{\frac{1}{3}}), \quad \text{if } \sigma(N, V) = \sigma(\mathcal{L}), \quad (1.34)$$

for some absolute constant C_0 .

Remark. Using this theorem we deduce that the function $h(u) = v(x(u)), u \in \mathbb{R}$ is continuous on $\mathbb{R} \setminus \{u_n, n \in \mathbb{Z}\}$, where $u_n = u(x), x \in g_n$. In this case we have

$$h(u_n \pm 0) \leq h(u_n), \quad n \in \mathbb{Z}. \quad (1.35)$$

The plan of our paper is as follows. In Sect. 2 we obtain the basic properties of the fundamental solution and prove Theorem 1.1. In Sect. 3 we determine the asymptotics of $M(z)$ and of the Lyapunov function and the multipliers at high energy and prove Theorem 1.2. Sect. 4 is the central part of the paper where we obtain the main properties of the quasimomentum $k_m, m = 1, \dots, N$. We prove the basic Theorems 4.1 and 4.2. In Sect. 5 using Theorems 4.1, 4.2 and [KK3], devoted to the conformal mapping theory, we prove Theorem 1.4 and 1.6.

2 Fundamental solutions

In this section we study ϑ, φ . We begin with some notational convention. A vector $h = \{h_n\}_1^N \in \mathbb{C}^N$ has the Euclidean norm $|h|^2 = \sum_1^N |h_n|^2$, while a $N \times N$ matrix A has the operator norm given by $|A| = \sup_{|h|=1} |Ah|$.

The fundamental solution φ satisfy the integral equations

$$\varphi(t, z) = \varphi_0(t, z) + \int_0^t \frac{\sin z(t-s)}{z} V(s) \varphi(s, z) ds, \quad \varphi_0(t, z) = \frac{\sin zt}{z} I_N, \quad (2.1)$$

where $(t, z) \in \mathbb{R} \times \mathbb{C}$. The standard iterations in (2.1) yields

$$\varphi(t, z) = \sum_{n \geq 0} \varphi_n(t, z), \quad \varphi_{n+1}(t, z) = \int_0^t \frac{\sin z(t-s)}{z} V(s) \varphi_n(s, z) ds. \quad (2.2)$$

The similar expansion $\vartheta = \sum_{n \geq 0} \vartheta_n$ with $\vartheta_0(t, z) = I_N \cos zt$ holds. In order to determine the asymptotics of the fundamental solutions we introduce the functions

$$I_m^0(z) = \int_0^m ds_1 \int_0^{s_1} \cos z(m - 2s_1 + 2s_2) F_2(s) ds_2, \quad m = 1, 2, \quad (2.3)$$

where $F_n(s) = \text{Tr } V(s_1) \cdot \dots \cdot V(s_n)$, $s = (s_1, \dots, s_n) \in \mathbb{R}^n$, $n \geq 1$. They satisfy the simple identity $I_2^0(z) = 4I_1^0(z) \cos z$, $z \in \mathbb{C}$, see [BKK]. Define

$$V^0 = \int_0^1 V(x) dx, \quad B_n = \text{Tr} \frac{(V^0)^n}{n!}, \quad n \geq 1, \quad |z|_1 = \max\{1, |z|\}. \quad (2.4)$$

We prove

Lemma 2.1. *Let $V \in \mathcal{H}$. Then each functions $\varphi(t, z), \vartheta(t, z), t \geq 0$, are entire and for any integers $m \geq t \geq 0, n_0 \geq -1$ the following estimates are fulfilled:*

$$\begin{aligned} \max \left\{ \left| \vartheta(t, z) - \sum_0^{n_0} \vartheta_n(t, z) \right|, \left| |z|_1 \left(\varphi(t, z) - \sum_0^{n_0} \varphi_n(t, z) \right) \right|, \left| \frac{1}{|z|_1} \left(\vartheta'(t, z) - \sum_0^{n_0} \vartheta'_n(t, z) \right) \right|, \right. \\ \left. \left| \varphi'(t, z) - \sum_0^{n_0} \varphi'_n(t, z) \right| \right\} \leq \frac{(m\kappa)^{n_0+1}}{(n_0+1)!} e^{t|\text{Im } z| + m\kappa}. \end{aligned} \quad (2.5)$$

Moreover, each $T_m(\cdot), m \geq 1$ is entire and satisfies

$$|T_m(z) - \sum_0^{n_0} T_m^{(n)}(z)| \leq \frac{(m\kappa)^{n_0+1}}{(n_0+1)!} e^{m(|\text{Im } z| + \kappa)}, \quad (2.6)$$

$$T_m^{(0)}(z) = \cos mz, \quad T_m^{(1)}(z) = \frac{mB_1}{2Nz} \sin mz, \quad T_m^{(2)}(z) = \frac{1}{4Nz^2} (I_2^{(m)}(z) - m^2 B_2 \cos mz), \dots$$

Proof. We prove the estimates of φ , the proof for $\varphi', \vartheta, \vartheta'$ is similar. (2.2) gives

$$\varphi_n(t, z) = \int_{D_n} f_n(t, s) V(s_1) \cdot \dots \cdot V(s_n) ds, \quad f_n(t, s) = \varphi_0(s_n, z) \prod_{k=1}^n \frac{\sin z(s_{k-1} - s_k)}{z}, \quad (2.7)$$

where $s = (s_1, \dots, s_n) \in \mathbb{R}^n$, $s_0 = t$ and $D_n = \{0 < s_n < \dots < s_2 < s_1 < t\}$. Substituting the estimate $|\varphi_0(t, z)| = |z^{-1} \sin zt| \leq |z|_1^{-1} e^{|\operatorname{Im} z|t}$ into (2.7), we obtain

$$|\varphi_n(t, z)| \leq \frac{e^{|\operatorname{Im} z|t}}{|z|_1^{n+1}} \int_{D_n} |V(s_1)| \cdot \dots \cdot |V(s_n)| ds \leq \frac{e^{|\operatorname{Im} z|t}}{|z|_1^{n+1}} \cdot \frac{1}{n!} \left(\int_0^t |V(x)| dx \right)^n.$$

This shows that for each $t \geq 0$ the series (2.2) converges uniformly on bounded subset of $\mathbb{C}$. Each term of this series is an entire function. Hence the sum is an entire function. Summing the majorants we obtain estimates (2.5).

The function $T_m, m \geq 1$ is entire, since φ, ϑ are entire. We have $(2N)T_m = \operatorname{Tr} M^m(z) = \operatorname{Tr} \mathcal{M}(m, z) = \operatorname{Tr} \sum_{n \geq 0} \mathcal{M}_n(m, z)$, where

$$\operatorname{Tr} \mathcal{M}_0(m, z) = 2N \cos mz, \quad \operatorname{Tr} \mathcal{M}_n(m, z) = \operatorname{Tr} \vartheta_n(m, z) + \operatorname{Tr} \varphi'_n(m, z), \quad n \geq 1. \quad (2.8)$$

The estimates $|\operatorname{Tr} \varphi'_n(m, z)| \leq \frac{(m\kappa)^n}{n!} e^{|\operatorname{Im} z|m}$ and $|\operatorname{Tr} \vartheta_n(m, z)| \leq \frac{(m\kappa)^n}{n!} e^{|\operatorname{Im} z|m}$ yield

$$|\operatorname{Tr} \mathcal{M}_n(m, z)| \leq (2N) \frac{(m\kappa)^n}{n!} e^{m|\operatorname{Im} z|}, \quad n \geq 0. \quad (2.9)$$

Using (2.8) we obtain

$$\operatorname{Tr} \mathcal{M}_1(m, z) = \frac{1}{z} \int_0^m (\sin z(m-t) \cos zt + \cos z(m-t) \sin zt) \operatorname{Tr} V(t) dt = \frac{\sin mz}{z} m B_1, \quad (2.10)$$

and

$$\begin{aligned} \operatorname{Tr} \mathcal{M}_2(m, z) &= \frac{1}{z^2} \int_0^m \int_0^t \sin z(t-s) z(m-t+s) F_2(t, s) dt ds \\ &= \frac{1}{2z^2} \int_0^m \int_0^t (\cos z(m-t+s) - \cos zm) F_2(t, s) dt ds = \frac{1}{2z^2} (I_m^0(z) - m^2 B_2 \cos mz) \end{aligned}$$

since $\int_0^m \int_0^t F_2(t, s) dt ds = \frac{1}{2} \operatorname{Tr} \left(\int_0^m V(t) dt \right)^2 = m^2 B_2$. ■

Note that the fundamental solutions $\varphi(t, z), \vartheta(x, z)$ and $M(z), T_m(z), m \geq 1$ is real for $z^2 \in \mathbb{R}$. Moreover, all functions $\varphi(t, z), \varphi'(t, z), \vartheta(t, z), \vartheta'(t, z), M(z), T_m, m \geq 1$ are even with respect to $z \in \mathbb{C}$ and then they are entire with respect to $\lambda = z^2$.

Using the identity (1.5) and $D(\tau, z) = \sum_{m=0}^{2N} \xi_m(z) \tau^{2N-m}$ we obtain

$$\frac{D(\tau, z)}{(2\tau)^N} = \frac{(\tau^N + \tau^{-N})}{2^N} + \xi_1 \frac{(\tau^{N-1} + \tau^{1-N})}{2^N} + \dots + \xi_{N-1} \frac{(\tau + \tau^{-1})}{2^N} + \dots + 2^{-N} \xi_N.$$

Substituting into this equality the identity for the Chebyshev polynomials $\mathcal{T}_n, n \geq 1$:

$$\frac{\tau^n + \tau^{-n}}{2} = \mathcal{T}_n(\nu) = 2^{N-1} \sum_0^{\lfloor \frac{n}{2} \rfloor} c_{n,m} \nu^{n-2m}, \quad c_{n,m} = (-1)^m n \frac{(n-m-1)!}{(n-2m)! m!} 2^{n-2m-N}, \quad (2.11)$$

$\nu = \frac{\tau + \tau^{-1}}{2}$, see [AS], we get $\Phi(\nu, z)$ given by

$$\Phi(\nu, z) = \frac{D(\tau, z)}{(2\tau)^N} = \sum_0^N \phi_m(z) \nu^{N-m}, \quad (2.12)$$

$$\phi_0 = 1, \quad \phi_1 = c_{N-1,0} \xi_1 = \frac{\xi_1}{2}, \quad \phi_2 = c_{N,1} + c_{N-2,0} \xi_2, \quad \phi_3 = c_{N-1,1} \xi_1 + c_{N-3,0} \xi_3, \dots, \quad (2.13)$$

$$\phi_{2n} = c_{N,n} + c_{N-2,n-1} \xi_2 + c_{N-4,n-2} \xi_4 + \dots + c_{N-2n,0} \xi_{2n}, \quad (2.14)$$

$$\phi_{2n+1} = c_{N-1,n} \xi_1 + c_{N-3,n-1} \xi_3 + c_{N-5,n-2} \xi_5 + \dots + c_{N-2n-1,0} \xi_{2n+1}. \quad (2.15)$$

Let D^0, Φ^0 be the determinant D and the polynomial Φ at $V = 0$. In this case we have

$$D^0(\tau, z) = (\tau^2 + 1 - 2\tau \cos z)^N = (2\tau)^N (\nu - \cos z)^N = (2\tau)^N \sum_0^N C_m^N (-\cos z)^m \nu^{N-m}$$

where $C_m^N = \frac{N!}{(N-m)!m!}$. Thus $\Phi^0 = (\nu - \cos z)^N$ and $\phi_m^0(z) = (-1)^m C_m^N \cos^m z$ at $V = 0$.

Substituting the identity for the Chebyshev polynomials (2.11) into the trace formula $\frac{1}{2} \text{Tr } M^n(z) = \sum_1^N \frac{\tau_m^n + \tau_m^{-n}}{2}$, we obtain

$$\sum_{m=1}^N \mathcal{T}_n(\Delta_m(z)) = \frac{1}{2} \text{Tr } M^n(z), \quad z \in \mathbb{C}. \quad (2.16)$$

Proof of Theorem 1.1 $\Delta_1(z), \dots, \Delta_N(z)$ are the roots of $\Phi(\nu, z) = 0$ for fixed $z \in \mathbb{C}$. Recall that $\mathcal{B}_\Delta$ are all branch points of $\Delta_1(z), \dots, \Delta_N(z)$.

First simple case, let $\mathcal{B}_\Delta = \emptyset$. Then all functions $\Delta_1(z), \dots, \Delta_N(z)$ are entire, the function $\Phi_j = \nu - \Delta_j, j = 1, \dots, N$ and $N_0 = N$.

Second case, let $\mathcal{B}_\Delta \neq \emptyset$. Consider a simply-connected domain $\Omega_1 \subset \mathbb{C}$ containing only one branch point $z_1 \in \mathcal{B}_\Delta$. We consider the behavior of the roots $\Delta_m, m = 1, \dots, N$ in the neighborhood of the branch points z_1 . Let $B'(z_1, r) = B(z_1, r) \setminus \{z_1\}$ be the small disk near z_1 with radius $r > 0$ but excluding z_1 . The functions $\Delta_m, m = 1, \dots, N$ are branches of analytic functions (defined in $\Omega_2 \setminus (\{z_1, z_2\})$) with a branch point (if $p_2 > 1$) at z_1 and z_2 .

are analytic of $z \in \mathcal{B}'(z_1, r)$. If $B'(z_1, r)$ is moved continuously around z_1 , then N functions can be continued analytically. When $B'(z_1, r)$ has been brought to its initial position after one revolution around z_1 , the functions $\Delta_m, m = 1, \dots, N$ will have undergone a permutation among themselves. These functions may therefore be grouped in the manner

$$\{\Delta_1(z), \dots, \Delta_{p_1}(z)\}, \{\Delta_{p_1+1}(z), \dots, \Delta_{p_1+q_1}(z)\}, \dots \quad \text{where } \Delta_i \neq \Delta_j, 1 \leq i < j \leq p_1, \dots \quad (2.17)$$

in such a way that each group undergoes a cyclic permutation by a revolution of $B'(z_1, r_1)$ of the kind described. For brevity each group will be called a cycle at the branch point z_1 , and the number of elements of a cycle will be called its period.

It is obvious that the elements of a cycle of period p_1 constitute a branch of an analytic function (defined near z_1) with a branch point (if $p_1 > 1$) at z_1 . We have Puiseux series as

$$\Delta_j(z) = \Delta_1(\zeta) + a_1 t + a_2 t^2 + \dots, \quad t = e^{j \frac{i2\pi}{p_1}} (z - z_1)^{\frac{1}{p_1}}, \quad j = 1, \dots, p_1.$$

Consider a simply-connected domain $\Omega_2 \subset \mathbb{C}$, containing only two distinct branch point $z_1, z_2 \in \mathcal{B}_\Delta$. The similar argument as above give that functions $\Delta_1, \dots, \Delta_N$ may therefore be grouped in the manner

$$\{\Delta_1(z), \dots, \Delta_{p_2}(z)\}, \{\Delta_{p_2+1}(z), \dots, \Delta_{p_2+q_2}(z)\}, \dots, \text{ where } p_1 \leq p_2, \dots \quad (2.18)$$

and $\Delta_i \neq \Delta_j, 1 \leq i < j \leq p_1, \dots$. Here the elements of a cycle of period p_2 constitute a branch of an analytic function (defined in $\Omega_2 \setminus (\{z_1, z_2\})$) with a branch point (if $p_2 > 1$) at z_1 and z_2 .

If we take a sequence of domain $\Omega_n \subset \Omega_{n+1}, n \geq 1$, containing only n distinct branch points $z_1, \dots, z_n \in \mathcal{B}_\Delta \cap \Omega_n$ and let $\lim \Omega_n = \mathbb{C}$, then the similar argument as above give that functions $\Delta_1, \dots, \Delta_N$ may therefore be grouped in the manner

$$\{\Delta_j(z), j \in \omega_1\}, \{\Delta_j(z), j \in \omega_2\}, \dots, \{\Delta_j(z), j \in \omega_{N_0}\}, \text{ where } \Delta_i \neq \Delta_j, i, j \in \omega_s, \quad (2.19)$$

$i \neq j, s = 1, \dots, N_0$. Here the elements of $\{\Delta_j(z), j \in \omega_1\}$ constitute a branch of an analytic function $\tilde{\Delta}_1$ (defined in $\mathbb{C} \setminus \mathcal{B}_\Delta$) with a branch point (if the of numbers of elements of ω_1 is > 1) at some points $z_n \in \mathcal{B}_\Delta, z \geq 1$.

There exists an interval $Y \subset \mathbb{R}, Y \neq \emptyset$ such that the spectral interval $Y \subset \sigma(\mathcal{L})$ has multiplicity $2N$ (see [MV]). Thus all functions $\Delta_m, m = 1, \dots, N$ are real on Y . Hence each entire function $\Phi_s(\nu, z), s = 1, \dots, N_0$ is real for all $\nu, z \in \mathbb{R}$.

ii) We have $\Delta'_m(z) = \frac{1}{2}(1 - \tau^{-2}(z))\tau'(z) \neq 0, z \in Y$, since by the Lyapunov Theorem, $\tau'(z) \neq 0$ for all $z \in Y$.

iii) Recall that the resultant for the polynomials $f = \tau^n + a_1\tau^{n-1} + \dots + a_n, g = b_0\tau^s + b_1\tau^{s-1} + \dots + b_s$ is given by

$$R(f, g) = \det \left(\begin{array}{cccccccc} 1 & a_1 & \dots & a_n & 0 & 0 & \dots & 0 \\ 0 & 1 & a_1 & \dots & a_n & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & 1 & a_1 & \dots & \dots & a_n \\ b_0 & b_1 & \dots & b_s & 0 & \dots & 0 & \dots \\ 0 & b_0 & b_1 & \dots & b_s & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & b_0 & b_1 & \dots & \dots & b_s \end{array} \right) \left. \begin{array}{l} \\ \\ \\ \\ \\ \\ \\ \end{array} \right\} \begin{array}{l} s \text{ lines} \\ \\ \\ n \text{ lines} \end{array} \quad (2.20)$$

The discriminant of the polynomial f with zeros $\tau_1, \dots, \tau_n$ is given by

$$\text{Dis} f = \prod_{i < j} (\tau_i - \tau_j)^2 = (-1)^{\frac{n(n-1)}{2}} R(f, f').$$

Thus we have $\text{Dis} \Phi_j(\tau, z) = \prod_{i < s, i, s \in \omega_j} (\Delta_i(z) - \Delta_s(z))^2 = (-1)^{\frac{N_j(N_j-1)}{2}} R(\Phi_j, \Phi'_j)$ is entire, since $\Phi_j(\tau, z)$ is the entire function. Then the function $\rho = \prod_1^{N_0} \text{Dis} \Phi_j$ is entire.

iv) We consider the gap $g_n = (z_n^-, z_n^+)$ in the variable z , where a gap $\gamma_n = (\lambda_n^-, \lambda_n^+), \lambda_n^\pm = (z_n^\pm)^2$. Each gap $g_n = (z_n^-, z_n^+) = \cup g_{n,p}, p = 1, \dots, p_n$, where $g_{n,p} = (z_{n,p}^-, z_{n,p}^+)$ is finite interval such that $\Delta_m(z) \notin [-1, 1]$ for all $z \in g_{n,p}$ for some $m = m(p)$. Note that $\Delta_m(z_{n,p}^-) = \pm 1$ or $z_{n,p}^-$ is the branch point $\in \mathcal{B}_\Delta$, otherwise we have a contradiction. ■

3 Spectral asymptotics

Below we need the identities for $J = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix}$, $J_1 = \begin{pmatrix} I_N & 0 \\ 0 & -I_N \end{pmatrix}$, $J_2 = \begin{pmatrix} 0 & I_N \\ I_N & 0 \end{pmatrix}$:

$$J^2 = -I, \quad J_1 J_2 = J, \quad J J_1 = -J_2, \quad J J_2 = J_1, \quad (3.1)$$

$$e^{zJ} = I_{2N} \cos z + J \sin z, \quad z \in \mathbb{C}. \quad (3.2)$$

We have also

$$D^0(z, \tau) = \det(e^{zJ} - \tau I_{2N}) = (\tau - e^{iz})^N (\tau - e^{-iz})^N, \quad D^0(z, \pm 1) = 2^N (1 \mp \cos z)^N, \quad (3.3)$$

$$(e^{zJ} - \tau I_{2N})^{-1} = (\tau - e^{iz})^{-1} (\tau - e^{-iz})^{-1} (e^{-zJ} - \tau I_{2N}). \quad (3.4)$$

We shall obtain the simple properties of the monodromy matrix. We introduce the modified monodromy matrix

$$\widetilde{M}(z) = \begin{pmatrix} I_N & 0 \\ 0 & zI_N \end{pmatrix}^{-1} M(z) \begin{pmatrix} I_N & 0 \\ 0 & zI_N \end{pmatrix} = \begin{pmatrix} \vartheta(1, z) & z\varphi(1, z) \\ z^{-1}\vartheta'(1, z) & \varphi'(1, z) \end{pmatrix}, \quad z \in \mathbb{C}, \quad (3.5)$$

with the same eigenvalues and the same traces. We will show following asymptotics

$$\widetilde{M}(z) = e^{zJ} \left(I + \frac{-V_0 J + \hat{V}(z) J_2}{2z} \right) + O(z^{-2} e^{|\operatorname{Im} z|}), \quad \hat{V}(z) = \int_0^1 V(t) e^{-2tzJ} dt, \quad |z| \rightarrow \infty. \quad (3.6)$$

Indeed, using (2.5), (3.5) we get $\widetilde{M}(z) = e^{zJ} + \widetilde{M}_1(z) + O(z^{-2} e^{|\operatorname{Im} z|})$, where $\widetilde{M}_1$ is given by

$$\widetilde{M}_1(z) = \frac{1}{2z} \int_0^1 V(t) f(t, z) dt, \quad f = 2 \begin{pmatrix} \sin z(1-t) \cos tz I_N & \sin z(1-t) \sin tz I_N \\ \cos z(1-t) \cos tz I_N & \cos z(1-t) \sin tz I_N \end{pmatrix}. \quad (3.7)$$

Let $c = \cos z$, $s = \sin z$ and $a = z(1-2t)$. Then substituting the identity

$$f = \begin{pmatrix} (s + \sin a) I_N & (-c + \cos a) I_N \\ (c + \cos a) I_N & (s - \sin a) I_N \end{pmatrix} = (s - cJ) + (\sin a J_1 - \cos a J_2) = -J e^{zJ} + e^{aJ} J_2$$

into (3.7) we obtain (3.6).

Define the matrix $L = \frac{1}{2}(\widetilde{M} + \widetilde{M}^{-1})$ with eigenvalues $\Delta_m = \frac{1}{2}(\tau_m + \tau_m^{-1})$, $m = 1, 2, \dots, N$, of multiplicity two. The identity (1.4) yields

$$L = \frac{\widetilde{M} + \widetilde{M}^{-1}}{2} = \frac{1}{2} \begin{pmatrix} \vartheta(1, \cdot) + \varphi'(1, \cdot)^\top & z(\varphi(1, \cdot) - \varphi(1, \cdot)^\top) \\ z^{-1}(\vartheta'(1, \cdot) - \vartheta'(1, \cdot)^\top) & \vartheta(1, \cdot)^\top + \varphi'(1, \cdot) \end{pmatrix}. \quad (3.8)$$

Using (2.5) we obtain

$$L(z) = L_1(z) + \frac{L_2(z)}{2z^2} + \frac{L_3(z)}{2z^3} + \frac{O(e^{|\operatorname{Im} z|})}{z^4}, \quad L_1(z) = \cos z + \frac{\sin z}{2z} V^0, \quad |z| \rightarrow \infty, \quad (3.9)$$

where

$$L_2(z) = \int_0^1 dt \int_0^t \sin z(t-s) \begin{pmatrix} a_{11}(t, s, z) & a_{12}(t, s, z) \\ a_{21}(t, s, z) & a_{22}(t, s, z) \end{pmatrix} ds, \quad z \in \mathbb{C}, \quad (3.10)$$

$$a_{11}(t, s, z) = \sin z(1-t) \cos zs V(t)V(s) + \cos z(1-t) \sin zs V(s)V(t), \quad a_{22} = a_{11}^\top, \quad (3.11)$$

$$a_{12}(t, s, z) = \sin z(1-t) \sin zs(V(t)V(s) - V(s)V(t)), \quad (3.12)$$

$$a_{21}(t, s, z) = \cos z(1-t) \cos zs(V(t)V(s) - V(s)V(t)), \quad (3.13)$$

and

$$L_3(z) = \int_0^1 dt \int_0^t ds \int_0^s \sin z(t-s) \sin z(s-u) \begin{pmatrix} b_{11}(t, s, u, z) & b_{12}(t, s, u, z) \\ b_{21}(t, s, u, z) & b_{22}(t, s, u, z) \end{pmatrix} du, \quad (3.14)$$

$$b_{11} = \sin z(1-t) \cos zu V(t)V(s)V(u) + \cos z(1-t) \sin zu V(u)V(s)V(t), \quad b_{22} = b_{11}^\top, \quad (3.15)$$

$$b_{12} = \sin z(1-t) \sin zu(V(t)V(s)V(u) - V(u)V(s)V(t)), \quad (3.16)$$

$$b_{21} = \cos z(1-t) \cos zu(V(t)V(s)V(u) - V(u)V(s)V(t)). \quad (3.17)$$

Recall that $V^0 = \int_0^1 V(x)dx$, $B_n = \text{Tr} \frac{(V^0)^n}{n!}$.

Lemma 3.1. *For each $(r, V) \in \mathbb{R}_+ \times \mathcal{H}$ asymptotics (1.11) and the following ones*

$$2^{2N} \det L(z) = \exp \left(-2Niz + i \frac{\text{Tr} V^0}{z} + \frac{i\|V\|^2 + o(1)}{4z^3} \right), \quad (3.18)$$

$$\text{Tr} L_2(z) = \left(-B_2 + \frac{i\|V\|^2 + o(1)}{2z} \right) \cos z, \quad (3.19)$$

$$\text{Tr} V_0 L_2(z) = -3B_3 \cos z + O(e^{|\text{Im} z|}/z), \quad (3.20)$$

$$\text{Tr} L_3(z) = -iB_3 \frac{\cos z}{2} + o(e^{|\text{Im} z|}), \quad (3.21)$$

hold as $y \geq r|x|, y \rightarrow \infty$. Moreover, the following identity and the asymptotics are fulfilled

$$L_2(\pi n) = \frac{(-1)^n}{4} \begin{pmatrix} X^{(n)} + [\hat{V}^{cn}, V^0] & [\hat{V}^{sn}, V^0 - \hat{V}^{cn}] \\ [\hat{V}^{sn}, V^0 + \hat{V}^{cn}] & X^{(n)} - [\hat{V}^{cn}, V^0] \end{pmatrix}, \quad (3.22)$$

$$\Delta_m(z) = \cos \left(z - \frac{V_m^0}{2z} \right) + \frac{O(|\hat{V}^{(n)}| + n^{-1})}{n^2} \quad \text{as } z = \pi n + O(1/n), \quad m = 1, \dots, N, \quad (3.23)$$

where $X^{(n)} = -(V^0)^2 + (\hat{V}^{cn})^2 + (\hat{V}^{sn})^2$ and $[A, B] = AB - BA$ for matrix A, B .

Proof. Recall the simple fact: Let A, B be matrices and $\sigma(B)$ be spectra of B . If A be normal, then $\text{dist}\{\sigma(A), \sigma(A+B)\} \leq |B|$ (see [Ka, p.291]).

The diagonal operator $L_1(z)$ has the eigenvalues $\Delta_m^0(z) = \cos z - V_m^0 \frac{\sin z}{2z}$, $m = 1, \dots, N$ with the multiplicity 2. Using the result from [Ka] and asymptotics (3.9) we deduce that the eigenvalues $\Delta_m(z)$ of matrix $L(z)$ satisfy the asymptotics (1.11).

Using (3.10) we obtain

$$\begin{aligned} \text{Tr } L_2(z) &= 2 \int_0^1 \int_0^t \sin z(t-s) \sin z(1-t+s) F_2(t, s) dt ds \\ &= \int_0^1 \int_0^t (\cos z(1-t+s) - \cos zm) F_2(t, s) dt ds = I_1^0(z) - B_2 \cos mz, \end{aligned}$$

since

$$\int_0^1 \int_0^t F_2(t, s) dt ds = \frac{1}{2} \text{Tr} \left(\int_0^1 V(t) dt \right)^2 = B_2.$$

Due to (6.1) we have $I_1^0(z) = \frac{i\|V\|^2 + o(1)}{2z} \cos z$, which yields (3.19).

We show (3.20). Let $G(t, s) = \text{Tr } V_0(V(t)V(s) + V(s)V(t))$. Using (3.10), (6.1) we have

$$\begin{aligned} \text{Tr } V_0 L_2(z) &= \text{Tr } V_0 \int_0^1 \int_0^t \sin z(t-s) z(1-t+s) (V(t)V(s) + V(s)V(t)) ds \\ &= \frac{1}{2} \int_0^1 dt \int_0^t (-\cos z + \cos z(1-t+s)) G(t, s) ds = -3B_3 \cos z + O(e^{|\text{Im } z|}/z), \end{aligned}$$

since

$$\int_0^1 dt \int_0^t \text{Tr } V_0(V(t)V(s) + V(s)V(t)) ds = \text{Tr } V_0 \left(\int_0^1 V(t) dt \right)^2 = 6B_3.$$

Consider $\text{Tr } L_3$. The identity (3.14) gives

$$\begin{aligned} \text{Tr } L_3(z) &= \int_0^1 dt \int_0^t ds \int_0^s \sin z(t-s) \sin z(s-u) \text{Tr}(b_{11} + b_{22}) du \\ &= \int_0^1 dt \int_0^t ds \int_0^s \sin z(t-s) \sin z(s-u) \sin z(1-t+u) R du, \end{aligned}$$

where $R = \text{Tr}(V(t)V(s)V(u) + V(u)V(s)V(t))$. Using the identity

$$4 \sin z(t-s) \sin z(s-u) \sin z(1-t+u) = -\sin z + P,$$

$$P = \sin z(1-2s+2u) - \sin z(1-2t+2s) - \sin z(1-2t+2u),$$

we get

$$L_3 = -L_3^0 \frac{\sin z}{2} + L_3^1, \quad L_3^0 = \frac{1}{2} \int_0^1 dt \int_0^t ds \int_0^s R du, \quad L_3^1 = \frac{1}{4} \int_0^1 dt \int_0^t ds \int_0^s P R du,$$

where

$$L_3^0 = \frac{\text{Tr}}{2} \int_0^1 V(t) dt \int_0^t ds \int_0^s (V(s)V(u) + V(u)V(s)) du = \frac{\text{Tr}}{2} \int_0^1 V(t) \left(\int_0^t V(s) ds \right)^2 = B_3.$$

Due to (6.3) we obtain $L_3^1 = o(e^{|\text{Im} z|})$, which yields (3.21).

We will show (3.18). Asymptotics (3.9) yields

$$\frac{L}{\cos z} = I_{2N} + S, \quad S = i \frac{V^0 I_{2N}}{2z} + \frac{L_2}{2z^2 \cos z} + \frac{L_3}{2z^3 \cos z} + O(z^{-4}), \quad (3.24)$$

as $|z| \rightarrow \infty, y \geq r|x|$, since $\tan z = i + O(e^{-y})$. In order to use the identity

$$\det(I + S) = \exp\left(\text{Tr} S - \text{Tr} \frac{S^2}{2} + \text{Tr} \frac{S^3}{3} + \dots\right), \quad |S| \rightarrow 0,$$

we need the traces of $S^m, m = 1, 2, 3$. Due to (3.19)-(3.22) we get

$$\frac{\text{Tr} S^3}{3} = -i \text{Tr} \frac{(V^0)^3 I_{2N}}{3(2z)^3} + O(z^{-4}) = -i \frac{B_3}{2z^3} + O(z^{-4}), \quad (3.25)$$

$$- \text{Tr} \frac{S^2}{2} = \frac{\text{Tr}}{2} \left(\frac{(V^0)^2}{(2z)^2} I_{2N} - 2i \frac{V^0 L_2}{4z^3 \cos z} + O(z^{-4}) \right) = \frac{B_2}{2z^2} + i \frac{3B_3}{4z^3} + O(z^{-4}) \quad (3.26)$$

and

$$\text{Tr} S = \text{Tr} \left(i \frac{V^0}{2z} + \frac{L_2}{2z^2 \cos z} + \frac{L_3 + O(z^{-1})}{2z^3 \cos z} \right) = i \frac{B_1}{z} + \left(-\frac{B_2}{2z^2} + i \frac{\|V\|^2}{4z^3} \right) - i \frac{B_3 + o(1)}{4z^3} \quad (3.27)$$

and summing (3.25)-(3.27) we get (3.18).

We will show (3.22). Let $z = \pi n, c_t = \cos \pi n t, s_t = \sin \pi n t$, and $K_{ts}^\pm = V(t)V(s) \pm V(s)V(t)$. Using (3.11), we have

$$\begin{aligned} 4(-1)^n \sin \pi n(t-s) a_{11}(t, s, \pi n) &= 4(s_t c_s - s_s c_t)(-s_t c_s V(t)V(s) + s_s c_t V(s)V(t)) \\ &= ((c_{2t} - 1)(1 + c_{2s}) + s_{2s} s_{2t}) V(t)V(s) + ((c_{2t} + 1)(1 - c_{2s}) + s_{2s} s_{2t}) V(s)V(t) \\ &= (-1 + c_{2t} c_{2s} + s_{2t} s_{2s}) K_{ts}^+ + ((c_{2t} - c_{2s}) K_{ts}^- \end{aligned}$$

and the integration yields

$$\int_0^1 dt \int_0^t \sin z(t-s) a_{11}(t, s, z) ds = \frac{1}{2} \int_0^1 \int_0^1 \sin z(t-s) a_{11}(t, s, z) dt ds = \frac{(-1)^n}{4} \left(X^{(n)} + [\hat{V}^{cn}, V^0] \right), \quad (3.28)$$

where $X^{(n)} = -(V^0)^2 + (\hat{V}^{cn})^2 + (\hat{V}^{sn})^2$. Using (3.12), we obtain

$$(-1)^n \sin \pi n(t-s) a_{12}(t, s, \pi n) = (s_t c_s - s_s c_t)(-s_t s_s) K_{ts}^- = (s_{2t} - s_{2s} + s_{2s} c_{2t} - c_{2s} s_{2t}) \frac{K_{ts}^-}{4}$$

and the integration yields

$$\int_0^1 dt \int_0^t \sin z(t-s) a_{12}(t, s, z) ds = \frac{(-1)^n}{4} (\hat{V}^{sn} V^0 - V^0 \hat{V}^{sn} - \hat{V}^{sn} \hat{V}^{cn} + \hat{V}^{cn} \hat{V}^{sn}). \quad (3.29)$$

The similar arguments give

$$\int_0^1 dt \int_0^t \sin z(t-s) a_{21}(t, s, z) ds = \frac{(-1)^n}{4} (V^0 \hat{V}^{sn} - \hat{V}^{sn} V^0 - \hat{V}^{sn} \hat{V}^{cn} + \hat{V}^{cn} \hat{V}^{sn}). \quad (3.30)$$

Combining the identities (3.28)-(3.30) and using $a_{22} = a_{11}^\top$ we obtain (3.22).

The diagonal operator $L_1(z)$ has the eigenvalues $\Delta_m^0(z) = \cos z - V_m^0 \frac{\sin z}{2z}$, $m = 1, \dots, N$ with the multiplicity 2. Using the result from [Ka] and asymptotics (3.22) we deduce that the eigenvalues $\Delta_m(z)$ of matrix $L(z)$ satisfy the asymptotics (3.23) as $z = \pi n + O(1/n)$. ■

Recall that $D^0(\pm 1, z) = (e^{iz} \mp 1)^N (e^{-iz} \mp 1)^N = 2^N (1 \mp \cos z)^N$.

Lemma 3.2. *Let $V \in \mathcal{H}$ and let $A = e^{|\operatorname{Im} z| + \varkappa}$, $\varkappa = \frac{\|V\|}{|z|_1}$, $|z|_1 = \max\{1, |z|\}$. Then*

i) The following estimates are fulfilled:

$$|\xi_m(z)| \leq (2NA)^m, \quad |\xi_m(z) - \xi_m^0(z)| \leq 2\varkappa(2NA)^m, \quad m = 1, \dots, N. \quad (3.31)$$

$$|D(\pm 1, z) - D^0(\pm 1, z)| < C_N \varkappa A^N, \quad C_N = 4(2N)^N. \quad (3.32)$$

ii) For each integer $n_0 > 4^N C_N \|V\|$ the function $D(1, z)$ has exactly $N(2n_0 + 1)$ roots, counted with multiplicity, in the disc $\{|z| < \pi(2n_0 + 1)\}$ and for each $|n| > n_0$, exactly $2N$ roots, counted with multiplicity, in the domain $\{|z - 2\pi n| < \frac{\pi}{2}\}$. There are no other roots.

iii) For each integer $n_0 > 4^N C_N \|V\|$ the function $D(-1, \lambda)$ has exactly $2Nn_0$ roots, counted with multiplicity, in the disc $\{|z| < 2\pi n_0\}$ and for each $|n| > n_0$, exactly $2N$ roots, counted with multiplicity, in the domain $\{|z - \pi(2n + 1)| < \frac{\pi}{2}\}$. There are no other roots.

iv) Assume that $V_i^0 \neq V_j^0$ for all $i \neq j \in \omega_s$ for some $s = 1, \dots, N_0$. Then there exists integer $n_0 \geq 1$ such that the function ρ_s has exactly $2N_s(N_s - 1)n_0$ roots, counted with multiplicity, in the disc $\{|z| < \pi(n_0 + \frac{1}{2})\}$ and for each $|n| > n_0$, exactly $N_s(N_s - 1)$ roots, counted with multiplicity, in the domain $\{|z - \pi n| < \frac{\pi}{2}\}$. There are no other roots.

Proof. We prove (3.31) by induction. Let the first estimate in (3.31) hold for ξ_m . Using (2.6) we obtain $|\xi_m(z)| \leq \mu^m$, $\mu = 2NA$ for $m = 0, 1, 2$. If $m \geq 2$, then substituting the estimate $|\xi_j(z)| \leq \mu^j$ and $|T_j(z)| \leq A^j$ (see (2.6)) into (1.6) we have

$$|\xi_{m+1}(z)| \leq \frac{2N}{m+1} \sum_0^m A^{m+1-j} \mu^j \leq \frac{2NA^{m+1}((2N)^{m+1} - 1)}{(m+1)(2N-1)} \leq \mu^{m+1}.$$

We shall show the second estimate in (3.31). Let $p_n = |\xi_n(z) - \xi_n^0(z)|$. The recurrent identities (1.6) and $|T_j(z) - T_j^{(0)}(z)| \leq \varkappa j A^j$, $|\xi_j(z)| \leq (2NA)^j$ and (2.6) give

$$p_n \leq \frac{2N}{n} \sum_0^{n-1} |T_{n-k}(z) \xi_k(z) - T_{n-k}^{(0)}(z) \xi_k^0(z)| \leq \frac{2N}{n} \sum_0^{n-1} (|\xi_k(z)| |T_{n-k}(z) - T_{n-k}^{(0)}(z)| + |T_{n-k}(z)| p_k)$$

$$\leq \frac{2N}{n} \sum_1^{n-1} \mu^k \varkappa(n-k) A^{n-k} + A^{n-k} \varkappa k \mu^k = 2N A^n \varkappa \sum_1^{n-1} (2N)^k \leq 2\varkappa \mu^n$$

where we used $p_1 \leq \varkappa A$, see (2.6). Thus we get (3.31).

ii) We have $D(1, \cdot) = \xi_N + 2 \sum_0^{N-1} \xi_n$, where ξ_n are given by (1.6). Recall that $\xi_n^0 = \xi_n$ at $V = 0$. Hence using (3.31) and $\mu = 2NA$ we have

$$|D(1, z) - D^0(1, z)| \leq |\xi_N(z) - \xi_N^0(z)| + 2 \sum_1^{N-1} |\xi_n(z) - \xi_n^0(z)| \leq 2\varkappa \mu^N + 2 \sum_1^{N-1} 2\varkappa \mu^n. \quad (3.33)$$

Thus $|D(1, z) - D^0(1, z)| < 2\varkappa \mu^N (1 + \frac{2}{\mu-1}) < 4\varkappa \mu^N$ since $\mu \geq 4$ which yields (3.32).

Let $n_1 > n_0$ be another integer. Introduce the contour $C_n(r) = \{z : |z - \pi n| = \pi r\}$. Consider the contours $C_0(2n_0 + 1), C_0(2n_1 + 1), C_{2n}(\frac{1}{4}), |n| > n_0$. Note that $\varkappa \leq \frac{1}{4^4}$ on all contours. Then (3.27) and the estimate $e^{\frac{1}{2}|\operatorname{Im} z|} < 4|\sin \frac{z}{2}|$ on all contours yield

$$\left| D(1, \lambda) - \left(2 \sin \frac{z}{2}\right)^{2N} \right| \leq C_N \varkappa A^N = C e^{N|\operatorname{Im} z|} \leq (C 4^N) \left| 2 \sin \frac{z}{2} \right|^{2N} < \frac{1}{2} \left| 2 \sin \frac{z}{2} \right|^{2N} \quad (3.34)$$

where $C = |z|^{-1} C_N \|V\| e^{N\varkappa}$. Hence, by Rouché's theorem, $D(1, z)$ has as many roots, counted with multiplicities, as $\sin^{2N} \frac{z}{2}$ in each of the bounded domains and the remaining unbounded domain. Since $\sin^{2N} \frac{z}{2}$ has exactly one root of the multiplicity $2N$ at $2\pi n$, and since $n_1 > n_0$ can be chosen arbitrarily large, the point ii) follows.

iii) The proof for $D(-1, \lambda)$ is similar.

iv) Consider the case $N_0 = 1$, the proof for $N_0 \geq 2$ is similar. Asymptotics (1.11) yields $\Delta_j(z) - \Delta_m(z) = (V_j^0 - V_m^0) \frac{\sin z}{2z} + O(z^{-2} e^{|\operatorname{Im} z|})$, $|z| \rightarrow \infty$. Then

$$\rho(z) = \prod_{j < m} (\Delta_j(z) - \Delta_m(z))^2 = c_0 \left(\frac{\sin z}{2z} + O\left(\frac{e^{|\operatorname{Im} z|}}{z^2}\right) \right)^{N(N-1)}, \quad c_0 = \prod_{j < m} (V_j^0 - V_m^0)^2. \quad (3.35)$$

Let $n_1 > n_0$ be another integer. Introduce the contour $C_n(r) = \{z : |z - \pi n| = \pi r\}$. Consider the contours $C_0(2n_0 + 1), C_0(2n_1 + 1), C_{2n}(\frac{1}{4}), |n| > n_0$. Note that $\varkappa \leq \frac{1}{4^4}$ on all contours. Then (3.35) and the estimate $e^{|\operatorname{Im} z|} < 4|\sin z|$ on all contours (for large n_0) yield

$$\rho(z) = \rho^0(z)(1 + O(z^{-1})), \quad \rho^0(z) = c_0 \left(\frac{\sin z}{2z} \right)^{N(N-1)}.$$

Hence, by Rouché's theorem, ρ has as many roots, counted with multiplicities, as ρ^0 in each of the bounded domains and the remaining unbounded domain. Since ρ^0 has exactly one root of the multiplicity $N(N-1)$ at $\pi n \neq 0$, and since $n_1 > n_0$ can be chosen arbitrarily large, the point iv) follows. ■

Proof of Theorem 1.2 i) We determine asymptotics (1.15) for $z_{m,m}^{n\pm} = \sqrt{\lambda_{m,m}^{n\pm}}$ as $n \rightarrow \infty$. Lemma 3.2 yields $|z_{m,m}^{n\pm} - \pi n| < \frac{\pi}{2}$ as $n \rightarrow \infty, m = 1, 2, \dots, N$. Lemma 3.1 gives $\Delta_m(z) = \cos(z - \frac{V_m^0}{2z}) + O(1/z^2)$ as $z = \pi n + O(1)$. Using the identity $\Delta_j(z_{m,m}^{n\pm}) = (-1)^n$, we have $z_{m,m}^{n\pm} = \pi n + O(1/n)$. We shall determine sharper asymptotics.

Recall that the modified monodromy matrix $\widetilde{M}(z)$ is given by (3.5) and it has the same eigenvalues as $M(z)$. Define the local parameter μ by $z = \pi n + \varepsilon\mu$, $\varepsilon = \frac{1}{2\pi n}$. Note that $\lambda = z^2 = (\pi n)^2 + \mu + (\varepsilon\mu)^2$. Asymptotics (3.6) gives

$$(-1)^n \widetilde{M}(z) = e^{\varepsilon\mu J} \left(I_{2N} - \varepsilon J(V_0 + J\hat{V}(\pi n)J_2) + O(\varepsilon) \right) = I_{2N} - \varepsilon J(V_0 + \hat{\mathcal{V}}^n + O(\varepsilon) - \mu), \quad (3.36)$$

where

$$\hat{\mathcal{V}}^n = J\hat{V}(\pi n)J_2 = J \int_0^1 V(t)e^{-2tzJ} dt J_2 = J(\hat{V}^{nc} - J\hat{V}^{ns})J_2 = J_1\hat{V}^{nc} + J_2\hat{V}^{ns}.$$

Hence we have the asymptotics (1.15), since

$$0 = \det \left((-1)^n \widetilde{M}(z) - I_{2N} \right) = \det(-\varepsilon J) \det \left(V_0 + \hat{\mathcal{V}}^n + O(\varepsilon) - \mu \right).$$

Consider the case $V_1^0 < \dots < V_N^0$. We shall determine asymptotics (1.18) for the case $\alpha = (m, m)$, $m = 1, \dots, N$. Let $z_{m,m}^{n\pm} = \pi n + \varepsilon\mu$ and $\mu - V_1^0 = \xi \rightarrow 0$. Using the simple transformation (unitary), i.e., changing the lines and columns, we obtain

$$\det(V^0 + \hat{\mathcal{V}}^{(n)} + O(\varepsilon) - \mu) = \det \begin{pmatrix} A_1 - \xi & A_2 \\ A_3 & A_4 - \xi \end{pmatrix} = \det(A_4 - \xi) \det K(\xi),$$

$$A_1 = \begin{pmatrix} \hat{V}_{11}^{cn} & \hat{V}_{11}^{sn} \\ \hat{V}_{11}^{sn} & -\hat{V}_{11}^{cn} \end{pmatrix} + O(\varepsilon), \quad A_4 = \text{diag}\{V_2^0 - V_1^0, \dots, V_N^0 - V_1^0\} + A_5,$$

$$A_2, A_3, A_5 = O(\delta_n), \quad \delta_n = |\hat{V}^n| + \varepsilon, \quad K(\xi) = A_1 - \xi - A_2(A_4 - \xi)^{-1}A_3 + O(\varepsilon)$$

We have $K(\xi) = A_1 - \xi + O(\phi)$, $\phi = \varepsilon + |\hat{V}^n|^2$, which yields

$$0 = \det K(\xi) = \xi^2 - |\hat{V}_{11}^n|^2 - \xi b_1 + \hat{V}_{11}^{cn} b_2 + \hat{V}_{11}^{sn} b_3 + O(\phi^2), \quad b_1, b_2, b_3 = O(\phi)$$

where b_1, b_2, b_3 are analytic functions of ξ . Rewriting the last equation in the form $(\xi + \alpha)^2 = (|\hat{V}_{11}^n| + \beta)^2 + O(\phi^2)$, $\alpha, \beta = O(\phi)$ and using the estimate $\sqrt{x^2 + y^2} - x \leq y$ for $x, y \geq 0$ we get $\xi = \pm |\hat{V}_{11}^n| + O(\phi)$, which yields (1.18) for the case $\alpha = (m, m)$.

Consider the resonances. Assume that $V_i^0 \neq V_j^0$ for all $i \neq j \in \omega_s$ for some $s = 1, \dots, N_0$. By Lemma 3.2, the zeros of ρ_s have the form $z_\alpha^{n\pm}$, $\alpha = (j, j')$, $j, j' \in \omega_s$, $j < j'$, $n \in \mathbb{Z} \setminus \{0\}$ and satisfy $|z_\alpha^{n\pm} - \pi n| < \pi/2$.

Asymptotics (1.11) yields $\Delta_j(z) - \Delta_{j'}(z) = (V_j^0 - V_{j'}^0) \frac{\sin z}{2z} + O(z^{-2}e^{|\text{Im } z|})$, $|z| \rightarrow \infty$. Then $|z_\alpha^{n\pm} - \pi n| < \pi/2$ yields $z_\alpha^{n\pm} = \pi n + O(1/n)$ as $n \rightarrow \infty$.

We have the identity $\Delta_j(z) - \Delta_{j'}(z) = 0$ at $z = z_\alpha^{n\pm}$. Then using (3.23) we have

$$\cos\left(z_\alpha^{n\pm} - \frac{V_j^0}{2\pi n}\right) - \cos\left(z_\alpha^{n\pm} - \frac{V_{j'}^0}{2\pi n}\right) = 2(-1)^n \sin \frac{V_{j'}^0 - V_j^0}{4\pi n} \sin\left(z_\alpha^{n\pm} - \pi n - \frac{V_j^0 + V_{j'}^0}{4\pi n}\right) = \frac{O(\delta_n)}{n^2}$$

which yields (1.16), i.e.,

$$z_\alpha^{n\pm} = \pi n + \varepsilon(a_\pm + \widetilde{z}_\alpha^{n\pm}), \quad a_\pm = \frac{V_j^0 \pm V_{j'}^0}{2}, \quad \varepsilon = \frac{1}{2\pi n}, \quad \text{wt } z_\alpha^{n\pm} = \delta_n. \quad (3.37)$$

We shall show that for large n in the neighborhood of each $\pi n + \varepsilon a_+$ the function $(\Delta_j(z) - \Delta_{j'}(z))^2$ has two real zeros resonances (counted with multiplicity). Introduce the functions

$$f_m(\mu) = 2(2\pi n)^2(1 - (-1)^n \Delta_m(\pi n + \varepsilon \mu)) = (\mu - V_m^0)^2 + O(\delta_n), \quad \delta_n = |\hat{V}^{(n)}| + \varepsilon. \quad (3.38)$$

For the case $\mu \rightarrow a_+$ we get

$$f_m(\mu) = (a_+ - V_m^0)^2 + o(1), \quad m = 1, \dots, N, \quad \text{and} \quad f_m(\mu) = a_-^2 + o(1), \quad m = j, j'. \quad (3.39)$$

Hence the function $f_j - f_{j'}$ (maybe) has the zeros, but the functions $f_j - f_m, m \neq j, j'$ have not zeros in the neighborhood of the point a_+ .

Note that these functions are real outside the small neighborhood of a_+ , otherwise for any complex branches there exists a complex conjugate branch, but the asymptotics (3.38) show that such branches are absent.

We have two cases: (1) let $f_m(\mu), m = j, j'$ be real in some small neighborhood of a_+ . Then the function $f_j - f_{j'}$ has at least one real zero, since by Theorem 1.1, the functions $f_j, f_{j'}$ are strongly monotone. Thus $(f_j - f_{j'})^2$ has at least 2 real zeros.

(2) Let $f_m(\mu), m = j, j'$ be complex in some small neighborhood of a_+ . Then they have at least two real branch points. Thus $(f_j - f_{j'})^2$ has at least 2 real zeros.

Hence $(f_j - f_{j'})^2$ has exactly two real zeros, since the number of resonances (in the neighborhood of the point πn) is equal to $N_s(N_s - 1)$.

We determine the sharp asymptotics of resonances. Define the unitary matrix $P = \frac{1}{\sqrt{2}}(J_1 + iJ) = P^*, P^2 = I_{2N}$. Using the identities

$$PJP = -iJ_1, \quad PJ_1P = iJ, \quad PJ_2P = -J_2,$$

$$\mathcal{V}_n = -iJ_1P\hat{V}^nP = -iJ_1(iJ\hat{V}^{cn} - J_2\hat{V}^{sn}) = J_2\hat{V}^{cn} + iJ\hat{V}^{sn} = \begin{pmatrix} 0 & \hat{V}^{(n)} \\ (\hat{V}^{(n)})^* & 0 \end{pmatrix}$$

we have

$$A = P\left(\frac{(-1)^n \widetilde{M}(z) - I}{i\varepsilon}\right)P = J_1(V^0 + P\hat{V}^nP + O(\varepsilon) - \mu) = J_1(V^0 - \mu) + i\mathcal{V}_n + O(\varepsilon).$$

The operator $A - a_-$ has the eigenvalue $\xi_0 = \frac{(-1)^n \tau_{n,s} - 1}{i\varepsilon} - a_-$ of multiplicity two, since $\tau_{n,s} = \tau(z_\alpha^{n\pm}) = (-1)^n e^{i\varepsilon(a_- + o(1))}$. The operator $J_1(V^0 - a_+) - a_- = \begin{pmatrix} V^0 - V_j^0 & 0 \\ 0 & -V^0 + V_{j'}^0 \end{pmatrix}$ has two eigenvalue ($= 0$) and other eigenvalues are not zeros. Using the simple transformation (unitary), i.e., changing the lines and columns, $\mu = a_+ + r \in \mathbb{R}, r = \widetilde{z}_\alpha^{n\pm} \rightarrow 0, v = \hat{V}_{jj'}^{(n)}$, we obtain

$$F(\xi) = \det(A - a_- - \xi) = \det \begin{pmatrix} A_1 - \xi & A_2 \\ A_4 & A_3 - \xi \end{pmatrix} = \det(A_3 - \xi) \det(K(\xi) - \xi I_2),$$

$$A_1 = \begin{pmatrix} -r & iv \\ i\bar{v} & r \end{pmatrix} + O(\varepsilon), \quad K(\xi) = A_1 - A_2(A_3 - \xi)^{-1}A_4 = \begin{pmatrix} -r + a_1 & iv + a_4 \\ i\bar{v} + a_3 & r + a_2 \end{pmatrix},$$

$A_3 = \text{diag}\{V_m^0 - V_j^0 - r, m \neq j\} \oplus \text{diag}\{V_{j'}^0 - V_m^0 + r, m \neq j'\} + A_4$, $A_2, A - 3, A_4 = O(\delta_n)$, the function $a_1, a_2, a_3, a_4 = O(\phi)$, $\phi = \varepsilon + |\hat{V}^{(n)}|^2$ and they analytic with respect to ξ in some small disk. The function $F = \det(K(\xi) - \xi I_2)$ has the form

$$F(\xi) = \xi^2 - r^2 + |v|^2 + a_1(r - \xi) + a_2(-r - \xi) - i v a_3 - i a_4 \bar{v} - a_4 a_3 = (\xi - \xi_0)^2 (1 + O(\xi - \xi_0)) \quad (3.40)$$

for $\xi \rightarrow 0$ where $\xi_0 = \frac{((-1)^n \tau_{n,s} - 1)}{i\varepsilon} - a_-$ is the zero of F of multiplicity two. Then $\xi_0 = O(\phi)$ and we have $(r - \alpha)^2 = (|v| - \beta)^2 + O(\phi^2)$ where $\alpha, \beta = O(\phi)$. Then using the estimate $\sqrt{x^2 + y^2} - x \leq y$ for $x, y \geq 0$ we get $r = \pm |v| + O(\phi)$. ■

Proof of Corollary 1.3 (i) Let $N_G = \infty$. Then, due to the Lyapunov Theorem, Theorem 1.2, there exists a real sequence $\lambda_k \rightarrow +\infty$ as $k \rightarrow \infty$, such that $\lambda_k \in \gamma_{n_k}^{\{j(m), m\}}$ for each $m = 1, \dots, N$. Hence, $\cap_{m=1}^N \gamma_{n_k}^{\{j(m), m\}} \neq \emptyset$. Using asymptotics (1.18) and $k \rightarrow \infty$, we obtain $V_1^0 + V_{j(1)}^0 = \dots = V_N^0 + V_{j(N)}^0$. Moreover, the estimates $V_1^0 < \dots < V_N^0$ yield $V_{j(1)}^0 > \dots > V_{j(N)}^0$, i.e. $j(1) = N, j(2) = N - 1, \dots$. Then, $V_1^0 + V_N^0 = V_2^0 + V_{N-1}^0 = \dots$, which give a contradiction.

(ii) Let $2a = V_1^0 + V_N^0 = V_2^0 + V_{N-1}^0 = \dots$. Due to (1.18), $(\pi n_k)^2 + a \in \cap_{m=1}^N \gamma_{n_k}^{\{N+1-m, m\}}$ as $k \rightarrow \infty$. Then the Lyapunov Theorem yields $(\pi n_k)^2 + a \notin \sigma(\mathcal{L})$, $k \rightarrow \infty$, i.e. $N_G = \infty$. ■

4 Harmonic functions

In this Sect. we will prove Theorems 4.1 and 4.2 about the properties of the quasimomentum. Recall that the Lyapunov function $\tilde{\Delta}_s(\zeta)$ is analytic on some N_s -sheeted Riemann surface $\mathcal{R}_s$ and $\mathcal{R} = \cup_1^{N_0} \mathcal{R}_s$. Let $z = x + iy \in \mathbb{C}$ be the natural projection of $\zeta \in \mathcal{R}$, $\mathcal{B}_\Delta$ be the set of all branch points of the Lyapunov function and $\mathcal{R}^\pm = \{\zeta \in \mathcal{R} : \pm \text{Im } \zeta > 0\}$. We define the simply connected domains $\mathcal{R}_0^\pm \subset \mathbb{C}_\pm$ and a domain $\mathcal{R}_0$ by

$$\mathcal{R}_0^\pm = \mathbb{C}_\pm \setminus \beta_\pm, \quad \beta_\pm = \bigcup_{\beta \in \mathcal{B}_\Delta \cap \mathbb{C}_\pm} [\beta, \beta \pm i\infty).$$

$$\mathcal{R}_0 = \mathbb{C} \setminus (\beta_+ \cup \beta_- \cup \beta_0), \quad \beta_0 = \{z \in \mathbb{R} : \Delta_m(z) \notin \mathbb{R} \text{ for some } m \in \{1, \dots, N\}\}$$

Due to the Lyapunov Theorem, $\Delta(\zeta) \notin [-1, 1]$, $\zeta \in \mathcal{R}^+$. Recall that $q(\zeta) = |\log \eta(\Delta(\zeta))|$ is the single-valued on $\mathcal{R}^+$ imaginary part of the (in general, many-valued on $\mathcal{R}^+$) quasimomentum $k(\zeta) = p(\zeta) + iq(\zeta) = \arccos \Delta(\zeta) = i \log \eta(\Delta(\zeta))$, where

$$\eta(z) \equiv z + \sqrt{z^2 - 1}, \quad \eta : \mathbb{C} \setminus [-1, 1] \rightarrow \{z \in \mathbb{C} : |z| > 1\}.$$

We denote by $q_m(z)$, $z \in \mathbb{C}_+$, $m = 1, \dots, N$, the branches of $q(\zeta)$ and by $p_m(z)$, $k_m(z)$, $z \in \mathcal{R}_0^+$, the single-valued branches of $p(\zeta)$, $k(\zeta)$, respectively.

Theorem 4.1. *Assume that $V \in \mathcal{H}$ and fix some $j = 1, \dots, N_0$. Then the function $\tilde{q}_s(\zeta) = \log |\eta(\tilde{\Delta}_s(\zeta))|$ is subharmonic on the Riemann surface $\mathcal{R}_s$ and the following asymptotics are fulfilled:*

$$\tilde{q}_s(\zeta) = y + O(1/|z|), \quad y > r|x|, \quad \text{any } r > 0, \quad (4.1)$$

$$\tilde{q}_s(\zeta) = y + O(|z|^{-\frac{1}{2}}), \quad (4.2)$$

as $|\zeta| \rightarrow \infty$, $\zeta \in \mathcal{R}_s$. Moreover, let Δ_j be analytic on some bounded interval $Y = (\alpha, \beta) \subset \mathbb{R}$ for some $j \in \omega_s$. Then

i) If $\Delta_j(z) \in \mathbb{R} \setminus [-1, 1]$ for all $z \in Y$, then $k_j(\cdot)$ has an analytic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$ such that

$$\operatorname{Re} k_j(z) = \operatorname{const} \in \pi\mathbb{Z}, \quad z \in Y, \quad (4.3)$$

$$q_j(z) = q_j(\bar{z}) > 0, \quad z \in \mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y. \quad (4.4)$$

ii) If we assume $\Delta_j(z) \notin \mathbb{R}$ for any $z \in Y$, then there exists a branch $\Delta_i, i \in \omega_s$ such that $\bar{\Delta}_i(z) = \Delta_j(z)$ for any $z \in Y$. The functions $\Delta_j(z)$ and $k_i + k_j$ have analytic extensions from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$ such that

$$\Delta_j(z) = \begin{cases} \Delta_j(z) & \text{if } z \in \mathcal{R}_0^+ \\ \bar{\Delta}_i(\bar{z}) & \text{if } z \in \mathcal{R}_0^- \end{cases}, \quad (4.5)$$

$$p_j(z) + p_i(z) = \operatorname{const} \in 2\pi\mathbb{Z}, \quad z \in Y, \quad q_j(z) = q_i(z), \quad z \in Y, \quad (4.6)$$

$$q_j(z) + q_i(z) = q_j(\bar{z}) + q_i(\bar{z}) > 0, \quad z \in \mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y. \quad (4.7)$$

Proof. By Theorem 1.1, the function $\tilde{\Delta}_s$ is analytic on $\mathcal{R}_s$ and the function $\eta(\cdot)$ is subharmonic on $\mathbb{C}$. Then $\tilde{q}_s(\zeta) = \log |\eta(\tilde{\Delta}_s(\zeta))|$ is subharmonic on $\mathcal{R}_s$. Using the asymptotics (1.20), (1.11) we obtain

$$\tilde{q}_s(\zeta) = \log |2 \cos z + O(y^{-1}e^y)| = y + O(y^{-1}), \quad |z| \rightarrow \infty, \quad y > r|x|,$$

which yields (4.1). Due to (1.11) and Lemma 6.2, we have

$$\tilde{q}_s(\zeta) = \log |\eta(\cos z + O(e^{|\operatorname{Im} z|}/z))| = \log |\eta(\cos z)| + O(1/\sqrt{z}) = y + O(1/\sqrt{z})$$

which yields (4.2).

i) Due to $\Delta_j(z) = \cos k_j(z)$ we obtain (4.3). The real part of k_j is a constant on Y , then k_j has an analytic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$. Moreover, q_j has an harmonic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$ by $q_j(z) = q_j(\bar{z})$, $z \in \mathcal{R}_0^\pm$.

ii) By Theorem 1.1, each polynomial $\Phi_s(\nu, z) = \prod_{n \in \omega_s} (\nu - \Delta_n(z))$, $z^2 \in \mathbb{R}$ is real for $\nu \in \mathbb{R}$. Then for Δ_j there exists a Δ_i such that $\Delta_j(x) = \bar{\Delta}_i(x)$, $x \in Y$. Then by the Morer Theorem, the function Δ_j has an analytic extension given by (4.5) from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$. Using $\Delta_m(z) = \cos k_m(z)$, $m = j, i$ we obtain $0 = \Delta_j(x) - \bar{\Delta}_i(x) = -2 \sin \frac{k_j(x) + \bar{k}_i(x)}{2} \sin \frac{k_j(x) - \bar{k}_i(x)}{2}$. Thus we get (4.6), (4.7) since $q_j(x) > 0$, $q_i(x) > 0$ on Y . ■

Recall the needed properties of the functions $v \in \mathcal{SC}$ defined in Sect. 1 and $w = u + iv$. It is well known, that $u \in C(\overline{\mathbb{C}}_+)$ and $\frac{1}{2\pi} \Delta v = \mu_v$ (in a sense of distribution) is a so-called Riesz measure of the function v . Moreover, the following identities are fulfilled:

$$\pi \mu_v((x_1, x_2)) = u(x_2) - u(x_1), \quad \text{for any } x_1 < x_2, \quad x_1, x_2 \in \mathbb{R}, \quad (4.8)$$

$$\frac{\partial v(z)}{\partial y} = y \int_{\mathbb{R}} \frac{d\mu_v(t)}{(t-x)^2 + y^2}, \quad z = x + iy \in \mathbb{C}_+, \quad (4.9)$$

which yields $\frac{\partial v(z)}{\partial y} \geq 0, z \in \mathbb{C}_+$. Moreover, $v(x) = v(x \pm i0), x \in \mathbb{R}$. It is well known that if $v \in \mathcal{SC}$, then

$$\int_{\mathbb{R}} \frac{d\mu_v(t)}{1+t^2} < +\infty, \quad \limsup_{z \rightarrow \infty} \frac{v(z)}{|z|} = \lim_{y \rightarrow +\infty} \frac{v(iy)}{iy} = \lim_{y \rightarrow +\infty} \frac{v(iy)}{y} = \lim_{x \rightarrow \pm\infty} \frac{u(x)}{x} \geq 0. \quad (4.10)$$

Now we recall the well known fact (see [Ah]).

Theorem (Nevanlinna). *i) Let μ be a Borel measure on $\mathbb{R}$ such that $\int_{\mathbb{R}} (1+x^{2p})d\mu(x) < +\infty$ for some $p \in \mathbb{Z}_+$. Then for each $r > 0$ the following asymptotics is valid*

$$\int_{\mathbb{R}} \frac{d\mu(t)}{t-z} = - \sum_{k=0}^{2p} \frac{Q_k}{z^{k+1}} + \frac{o(1)}{z^{2p+1}}, \quad |z| \rightarrow \infty, \quad y > r|x|, \quad Q_n = \int_{\mathbb{R}} x^n d\mu(x), \quad 0 \leq n \leq 2p.$$

ii) Let F be an analytic function in $\mathbb{C}_+$ such that $\text{Im } F(z) \geq 0, z \in \mathbb{C}_+$ and

$$\text{Im } F(iy) = c_0 y^{-1} + \dots + c_{2p-1} y^{-2p} + O(y^{-2p-1}) \quad \text{as } y \rightarrow \infty \quad (4.11)$$

for some $c_0, \dots, c_{2p-1}$ and $p \geq 0$. Then $F(z) = C + \int_{\mathbb{R}} \frac{d\mu(t)}{t-z}, z \in \mathbb{C}_+$, for some Borel measure μ on $\mathbb{R}$ such that $\int_{\mathbb{R}} (1+x^{2p})d\mu(x) < \infty$ and $C \in \mathbb{R}$.

Theorem 4.2. Assume that $V \in \mathcal{H}$. Then the function $w = \frac{1}{N} \sum_1^N k_m$ is analytic in $\mathbb{C}_+$; the function $v = \text{Im } w = \frac{1}{N} \sum_1^N q_m$ belongs to $\mathcal{SK}_0^+ \cap C(\mathbb{C})$ and it is positive harmonic in $\Omega = \mathbb{C}_+ \cup \mathbb{C}_- \cup g$. Moreover, the following asymptotics and identities hold

$$u(z) = \text{const} \in \frac{\pi}{N} \mathbb{Z}, \quad v(z) > 0, \quad z \in g_n, \quad (4.12)$$

$$v(z) = y + O(1/y), \quad |z| \rightarrow \infty, \quad y > r|x|, \quad \text{any } r > 0, \quad (4.13)$$

$$v(z) = y + O(|z|^{-\frac{1}{2}}) \quad |z| \rightarrow \infty. \quad (4.14)$$

Proof. Recall that $k_m(z) = i \log \eta(\Delta_m(z)), z \in \mathcal{R}_0^+$, where $\Delta_m(z), m = 1, \dots, N$, are the branches of the function Δ analytic on $\mathcal{R}$. Since $\Delta(\zeta) \notin [-1, 1], \zeta \in \mathcal{R}^+$, the function $\eta(\Delta(\zeta))$ is analytic on $\mathcal{R}^+$ and $|\eta(\Delta(\zeta))| > 1, \zeta \in \mathcal{R}^+$. Recall that $\mathcal{B}_\Delta$ is the set of all branch points of the function Δ . Define the function $F(z) \equiv \prod_{m=1}^N \eta(\Delta_m(z)), z \in \mathbb{C}_+$. Due to the symmetry of F with respect to the permutations of $\Delta_1, \dots, \Delta_n$, we deduce that F is analytic in the domain $B(z_0, r) \setminus \{z_0\}$ for any $z_0 \in \mathbb{C}_+$ and some $r = r(z_0)$, including the case $z_0 \in \mathbb{C}_+ \cap \mathcal{B}_\Delta$. Note that F is bounded in $B(z_0, r)$. This implies F is analytic in $B(z_0, r)$ for any $z_0 \in \mathbb{C}_+$, i.e. F is analytic in $\mathbb{C}_+$ and $|F(z)| > 1, z \in \mathbb{C}_+$. Therefore, the averaged quasimomentum

$$w(z) = \frac{i}{N} \log F(z) = \frac{1}{N} \sum_{m=1}^N i \log \eta(\Delta_m(z)) = \frac{1}{N} \sum_{m=1}^N k_m(z) \quad (4.15)$$

is analytic in $\mathbb{C}_+$ and $v(z) = \text{Im } w(z) > 0, z \in \mathbb{C}_+$. Moreover, w is continuous in $\overline{\mathbb{C}_+}$. Using this fact and Theorem 4.1 we deduce that v has an harmonic extension from $\mathbb{C}_+$ into Ω by

$v(z) = v(\bar{z})$, $z \in \mathbb{C}_\pm$. Moreover, w has an analytic extension from $\mathbb{C}_+$ into $\mathbb{C}_+ \cup \mathbb{C}_- \cup g_n$ since $u = \text{const}$ on the gap g_n , and we have identities (4.12).

We show that v is subharmonic in $\mathbb{C}$. Since v is harmonic in $\mathbb{C}_+ \cup \mathbb{C}_- \cup g$ we have to show that v is subharmonic at $z_0 \in \mathbb{R} \setminus g$. Note that v is continuous in $\mathbb{C}$.

Introduce the sets

$$\alpha_s(z) = \{m : \Delta_m(z) \in [-1, 1]\}, \quad \alpha_g(z) = \{m : \Delta_m(z) \in \mathbb{R} \setminus [-1, 1]\},$$

and $\alpha_c(z) = \{m : \Delta_m(z) \notin \mathbb{R}\}$. We have the decomposition

$$v = v_s + v_c + v_g, \quad v_j = \sum_{m \in \alpha_j(z_0)} \frac{q_m}{N}, \quad j = s, c, g.$$

We have 2 cases: the first case $z_0 \notin \mathcal{B}_\Delta$.

Consider v_g . We take the interval $Y = (z_0 - \varepsilon, z_0 + \varepsilon)$ for some $\varepsilon > 0$ such that $Y \cap \mathcal{B}_\Delta = \emptyset$. Due to Theorem 4.1 each $q_m(\cdot)$, $m \in \alpha_g(z_0)$ has an harmonic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$ by $q_m(z) = q_m(\bar{z})$, $z \in \mathcal{R}_0^\pm$ for sufficiently small ε . Thus v_g has an harmonic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$ by $v_g(z) = v_g(\bar{z})$, $z \in \mathcal{R}_0^\pm$.

Consider v_c . Due to Theorem 4.1 we deduce that

$$v_c = \sum_{m \in \alpha_c(z_0)} q_m(z) = \frac{1}{2} \sum_{m \in \alpha_c(z_0)} q_m(z) + q_{\tilde{m}}(z),$$

where $k_{\tilde{m}}(z)$, $\tilde{m} \in \alpha_c$, is the quasimomentum such that $\Delta_{\tilde{m}}(z) = \bar{\Delta}_m(z)$, $z \in Y$. Due to Theorem 4.1 each $v_m = q_m(\cdot) + q_{\tilde{m}}$, $m \in \alpha_g(z_0)$ has an harmonic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$ by $v_m(z) = v_m(\bar{z})$, $z \in \mathcal{R}_0^\pm$ for sufficiently small ε . Thus v_c has an harmonic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y$ by $v_c(z) = v_c(\bar{z})$, $z \in \mathcal{R}_0^\pm$.

Hence due to $v_s(z_0) = 0$ we obtain $v(z_0) \leq \frac{1}{2\pi} \int_0^{2\pi} v(z_0 + re^{i\phi}) d\phi$, for any small $r > 0$, and thus v is subharmonic in some neighborhood of z_0 .

The second case $z_0 \in \mathcal{B}_\Delta$. Define two intervals $Y_- = (z_0 - 2\varepsilon, z_0)$, $Y_+ = (z_0, z_0 + 2\varepsilon)$, for some $\varepsilon > 0$ such that $Y_\pm \cap \mathcal{B}_\Delta = \emptyset$ and each q_m is harmonic in $\{|z - z_0| < \varepsilon\} \cap \mathbb{C}_+$. Define the functions

$$v = v_1 + v_0, \quad v_0(z) = \sum_{m \in \alpha_0} q_m(z), \quad \alpha_0 = \alpha_s(z_0 + \varepsilon) \cup \alpha_g(z_0 + \varepsilon).$$

Note that $v_0(z_0) = 0$. The above arguments show that the function v_1 has a harmonic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup Y_- \cup Y_+$ by $v_1(z) = v_1(\bar{z})$ for sufficiently small $\varepsilon > 0$. Then the function v_1 has an harmonic extension from $\mathcal{R}_0^+$ into $\mathcal{R}_0^+ \cup \mathcal{R}_0^- \cup (z_0 - 2\varepsilon, z_0 + 2\varepsilon)$ by $v_1(z) = v_1(\bar{z})$ for sufficiently small ε , since the function v_1 is bounded in the disk $B(z_0, t) = \{z : |z - z_0| < \varepsilon\}$ for some small $\varepsilon > 0$. Thus we obtain $v(z_0) \leq \frac{1}{2\pi} \int_0^{2\pi} v(z_0 + re^{i\phi}) d\phi$ for any small $r > 0$, since $v_0(z_0) = 0$. Hence the function v is subharmonic in $\mathbb{C}$.

Asymptotics (4.1), (4.2) give (4.12), (4.13). Moreover, (4.13) yields $\int_{\mathbb{R}} \frac{v(t)}{1+t^2} dt < \infty$ and $\limsup_{z \rightarrow \infty} \frac{v(z)}{|z|} = 1$, i.e., $v \in \mathcal{SC}$.

The function $v(z) - y$ is harmonic in $\mathbb{C}_+$, it is positive on the real line and $v(z) - y = o(1)$ as $|z| \rightarrow \infty$, $z \in \overline{\mathbb{C}}_+$. Then $v(z) - y$ is positive on $\mathbb{C}_+$. Using (4.12) and the Nevanlinna Theorem we deduce that $\int_{\mathbb{R}} v(t) dt < \infty$. Then $v \in \mathcal{SK}_0^+$. ■

5 The conformal mappings

We need the result from [KK1].

Theorem 5.1. *Let $v \in \mathcal{SK}_0^+$ and $v \neq \text{const}$. Then $w : \mathbb{C}_+ \rightarrow w(\mathbb{C}_+) = W(h)$ is a conformal mapping for some $h \in C_{us}, h \geq 0$. Moreover, the following asymptotics, estimates and identities are fulfilled:*

$$w(z) = z - \frac{Q_0 + o(1)}{z}, \quad |z| \rightarrow \infty, \quad y \geq r|x|, \quad \text{for any } r > 0, \quad (5.1)$$

$$I_0^D + \mathcal{P}_0 = Q_0, \quad \sup_{x \in \mathbb{R}} v^2(x) \leq 2Q_0. \quad (5.2)$$

If in addition, $Q_2 < \infty$, then $Q_2 = I_1^D + \mathcal{P}_2$, where $\mathcal{P}_n, I_n^D, Q_n, n \geq 0$ are given by (1.23).

Proof of Theorem 1.4. By Theorem 4.2, the function $v = \frac{1}{N} \sum_1^N q_m \in \mathcal{SK}_0^+$. Then Theorem 5.1 gives that $w : \mathbb{C}_+ \rightarrow w(\mathbb{C}_+) = W(h)$ is a conformal mapping for some $h \in C_{us}$.

The function $M(iy), y > 0$ is real, then $M(z) = \overline{M}(-\bar{z}), z \in \overline{\mathbb{C}}_+$. This yields that the set $\{\tau_m(z)\}_1^N = \{\tau_m(-\bar{z})\}_1^N, z \in \overline{\mathbb{C}}_+$, which gives $v(z) = v(-\bar{z}), z \in \overline{\mathbb{C}}_+$. Thus $v(x) = v(-x)$ for all $x \in \mathbb{R}$ and the identities

$$w(-\bar{z}) = -\bar{z} + \frac{1}{\pi} \int_{\mathbb{R}} \frac{v(t)dt}{t + \bar{z}} = -\bar{z} - \frac{1}{\pi} \int_{\mathbb{R}} \frac{v(s)ds}{s - \bar{z}} = -\overline{w}(z), z \in \mathbb{C}_+$$

give $-w_0(-\bar{z}) = \overline{w}_0(z), z \in \mathbb{C}_+$.

Using (1.11) and $q_m \geq 0$ we obtain $q_m(iy) = y + o(1)$ and

$$\det L(z) = \prod_1^N \Delta_m^2(z) = \frac{(1 + e^{-4Ny}O(1))}{4^N} e^{-2i \sum_1^N k_m(z)} = \frac{(1 + e^{-4Ny}O(1))}{4^N} e^{-2iNw(z)} \quad (5.3)$$

as $z = iy, y \rightarrow \infty$ and (3.18) yields

$$\det L(z) = (\cos^{2N} z) \exp\left(i \frac{\text{Tr } V^0}{z} + \frac{i\|V\|^2 + o(1)}{4z^3}\right). \quad (5.4)$$

Thus due to $\text{Re } w(iy) = 0$ we get

$$w(z) = z - \frac{\text{Tr } V^0}{2Nz} - \frac{\|V\|^2 + o(1)}{8Nz^3}, \quad z = iy, \quad y \rightarrow \infty. \quad (5.5)$$

Then by the Nevanlinna Theorem and Theorem 5.1, we have asymptotics (1.25) and identities (1.26), (1.27).

Estimate (5.2) gives $v|_{\mathbb{R}} \leq \sqrt{2Q_0}$. If $z \in \sigma_{(N)}$, then $v(z) = 0$, since $\Delta_m(z) \in [-1, 1]$ and $q_m(z) = 0$ for all $m = 1, \dots, N$. If $z \in \sigma_{(1)} \cup g$, then $\Delta_m(z) \notin [-1, 1]$ for some $m = 1, \dots, N$. Thus $q_m(z) > 0$ and $v(z) > 0$, which gives (1.28). ■

Proof of Theorem 1.6. i) We show (1.30). Using $w(z) = z + \frac{1}{\pi} \int_{\mathbb{R}} \frac{v(t)dt}{(t-x)^2}, z \in \mathbb{C}_+$ and (1.28) we obtain $w'(z) = 1 + \frac{1}{\pi} \int_{\mathbb{R}} \frac{v(t)dt}{(t-x)^2} > 1, z \in \sigma_{(N)}$, which yields $u'_x(z) > 1, z \in \sigma_{(N)}$. We used the fact that due to (1.17) $u'_x(z) = 1$ for some $z \in \sigma_{(N)}$ iff $V = 0$.

Let $x \in \sigma_{(1)}$. Then some branch $\Delta_m(x) \in (-1, 1)$ for all $x \in Y \subset \sigma_{(1)}$ for some small interval $Y = (\alpha, \beta)$. We have $\Delta_m(x) = \cos k_m(x)$. Thus we get $k'_m(x) = -\frac{\Delta'_m(x)}{\sin k_m(x)} \neq 0$. Hence we get $u'_m(x) = k'_m(x) > 0$, since $q_m(x) = 0$ and (4.9) yields $u_x(x) \geq 0$ for $x \in Y$, which gives (1.30).

We show (1.31). Note that $v(z) > 0, z \in g$, otherwise we have not a gap. Using (4.9) we obtain $-\frac{\partial^2 v(z)}{\partial x^2} = \frac{\partial^2 v(z)}{\partial y^2} = \int_{\mathbb{R}} \frac{d\mu_v(t)}{(t-x)^2} > 0$, $z = x \in g_n$, which yields $v''_{xx}(z) < 0, z \in g_n$. Consider the function $u(z), z \in g_n$. Theorem 4.1 yields $N \operatorname{Re} w(z) = \sum_1^N \operatorname{Re} k_m = \sum_1^N \pi n_m = \pi N_n$, which gives (1.31).

We recall the result from [KK2]. Let a function f be harmonic and positive in the domain $\mathbb{C} \setminus [-a, a]$, $a > 0$ and $f(iy) = y(1 + o(1))$ as $y \rightarrow \infty$. Assume $f(z) = f(\bar{z})$, $z \in \mathbb{C} \setminus [-a, a]$ and $f \in C(\overline{\mathbb{C}_+})$. Then

$$f(x) = \sqrt{a^2 - x^2} \left(1 + \frac{1}{\pi} \int_{\mathbb{R} \setminus I} \frac{f(t) dt}{|t - x| |t^2 - a^2|^{\frac{1}{2}}} \right), \quad x \in [-a, a].$$

Hence the last identity and properties of v yield (1.32) and the estimate $v(z) \geq v_n^0(z) = |(z - z_n^-)(z_n^+ - z)|^{\frac{1}{2}}$, $z \in g_n = (z_n^-, z_n^+)$. Consider the case $g_n \subset \mathbb{R}_+$ (the proof for the cases $g_n \subset \mathbb{R}_-$ is similar). Defining $z_n^0 = \frac{z_n^+ + z_n^-}{2}$, $r = \frac{|g_n|}{2}$ and using $(z_0 + x)^{2p} + (z_0 - x)^{2p} \geq 2z_0^{2p}$ for all $p \geq 0$ we have

$$\frac{1}{\pi} \int_{g_n} t^{2p} v(t) dt \geq \int_{g_n} \frac{t^{2p} v_n^0(t) dt}{\pi} \geq \frac{1}{\pi} \int_0^r \sqrt{r^2 - x^2} ((z_n^0 + x)^{2p} + (z_n^0 - x)^{2p}) dx \geq \frac{r^2}{2} (z_n^0)^{2p}.$$

If $p = 0$, then we get $Q_0 = \frac{1}{\pi} \int_{\mathbb{R}} v(t) dt \geq \frac{1}{8} \sum |g_n|^2$, which yields the first estimate in (1.33).

If $p = 1$, then using $\gamma_n = \lambda_n^+ - \lambda_n^- \geq 0$, $\lambda_n^\pm = (z_n^\pm)^2$ we get

$$Q_2 = \frac{1}{\pi} \int_{\mathbb{R}} t^2 v(t) dt \geq \frac{1}{8} \sum_{n \in \mathbb{Z}} |g_n|^2 \frac{(z_n^+ + z_n^-)^2}{4} = \frac{1}{16} \sum_{n \geq 0} |\gamma_n|^2,$$

which yields the last inequality in (1.33).

Recall estimates from [K10]. The conformal mapping $w : \mathbb{C}_+ \rightarrow W(h)$ for the case $w(z) = z + o(1)$ as $|z| \rightarrow \infty$ and $h(u) = 0, u \neq \frac{\pi}{N}\mathbb{Z}$ and $\{h(\frac{\pi}{N}n)\}_{-\infty}^\infty \in \ell_1^2$ was studied in [K7]. For some absolute constant C_0 the following estimate was obtained: $Q_2 \leq C_0 G^2 (1 + G^{\frac{1}{3}})$, which together with $Q_2 = \frac{\|V\|^2}{8N}$ gives the estimate (1.34). ■

6 Appendix

Lemma 6.1. *Let function $F(s, t), R(t, s, u), t, s, u \in (0, 1)$ satisfy :*

- i) $F(\cdot, \cdot) \in L^2((0, 1)^2)$ and $F(t, t) \in L^1(0, 1)$,
- ii) $R(\cdot, \cdot, \cdot) \in L^2((0, 1)^3)$ and $R(t, t, s), R(t, s, t), R(s, t, t) \in L^1((0, 1)^2)$. Then

$$f(z) = \int_0^1 dt \int_0^t \cos z (1 - 2t + 2s) F(t, s) ds = \frac{i \cos z}{2z} \left(\int_0^1 F(t, t) dt + o(1) \right) \quad (6.1)$$

$$f^+(z) = \int_0^1 dt \int_0^t e^{-i2z(t-s)} F(t, s) ds = \frac{o(e^{2y})}{|z|}, \quad (6.2)$$

$$\int_0^1 dt \int_0^t ds \int_0^s \sin z(1-2\zeta) R(t, s, u) du = o(e^y), \quad (6.3)$$

as $r|x| < y \rightarrow \infty$ for any fixed $r > 0$, where ζ is one of functions: $s-u, t-u$ or $t-s$.

Proof. We have $f(z) = e^{iz} f^+(z) + e^{-iz} f^-(z)$, where

$$f^-(z) \equiv \frac{1}{2} \int_0^1 dt \int_0^t e^{i2z(t-s)} F(t, s) ds = \frac{1}{4} \int_0^1 ds \int_0^1 e^{i2z|t-s|} F(t, s) ds \quad (6.4)$$

where $F(t, s) = F(s, t), t, s \in (0, 1)$. Substituting the identities

$$\frac{1}{2\pi} \int_{\mathbb{R}} e^{ipk} \frac{dk}{k^2 - 4z^2} = \frac{i}{4z} e^{i2z|p|}, \quad p \in \mathbb{R}, \quad z \in \mathbb{C}_+, \quad \hat{F}(r, k) \equiv \frac{1}{2\pi} \iint_{[0,1]^2} e^{-irt-iks} F(t, s) dt ds,$$

into (6.4) we obtain

$$f^- = \frac{z}{i} \int_{\mathbb{R}} \frac{\hat{F}(-k, k) dk}{k^2 - 4z^2} = \frac{1}{i4z} \int_{\mathbb{R}} \left(-1 + \frac{k^2}{k^2 - 4z^2} \right) \hat{F}(-k, k) dk = \frac{-1}{i4z} \left(\int_0^1 F(t, t) dt + o(1) \right).$$

Consider f^+ . Let $\|F\|^2 = \int_0^1 dt \int_0^t |F(t, s)|^2 ds$. We have

$$|f^+(z)|^2 \leq g(z) \|F\|^2, \quad g(z) = \int_0^1 dt \int_0^t e^{4y(t-s)} ds \leq \int_0^1 \frac{e^{4yt}}{4y} dt \leq \frac{e^{4y}}{(4y)^2}. \quad (6.5)$$

Let F_0 be a smooth function such that $\|F - F_0\| \leq \varepsilon$ for some small $\varepsilon > 0$. Define the function $f_0^+(z) = \int_0^1 dt \int_0^t e^{-i2z(t-s)} F_0(t, s) ds$. Using (6.5) we obtain

$$|f^+(z)| \leq |f_0^+(z)| + |f^+(z) - f_0^+(z)| \leq |f_0^+(z)| + \|F - F_0\| \frac{e^{2y}}{(4y)} \quad (6.6)$$

and the integration by parts yields $f_0^+(z) = O(\frac{e^{2y}}{y^2})$. Thus we obtain (6.2).

The proof for (6.3) is similar. ■

Lemma 6.2. *The function $f(z) = \log |\xi(z)|, z \in \mathbb{C} \setminus [-1, 1]$ is subharmonic and continuous in $\mathbb{C}$. Moreover, for some absolute constant C the following estimate is fulfilled:*

$$|f(z) - f(z_0)| \leq C\varepsilon^{\frac{1}{2}}, \quad \text{if } |z - z_0| \leq \varepsilon \max\{2, |z_0|\}, \quad 0 \leq \varepsilon \leq \frac{1}{8}, \quad z, z_0 \in \mathbb{C}. \quad (6.7)$$

Proof. The function $f(z) = \log |\xi(z)|, z \in \mathbb{C} \setminus [-1, 1]$ is harmonic and $f(z) = 0, z \in [-1, 1]$. Then f is subharmonic in $\mathbb{C}$.

1. Let $|z_0| \leq 2$. Consider the case $|z_0 - 1| \leq 1/2$. The proof of other cases is similar. Firstly, let $z, z_0 \in \overline{\mathbb{C}}_+$. Then

$$f(z) - f(z_0) = \operatorname{Re} \int_{z_0}^z \frac{dt}{\sqrt{t^2 - 1}} = \operatorname{Re} \int_{z_0-1}^{z-1} \frac{ds}{\sqrt{s(2+s)}}, \quad \frac{1}{\sqrt{s(2+s)}} = \frac{1}{\sqrt{s}} + H(s),$$

where $H(\cdot)$ is analytic and bounded in the disk $B(0, 1)$. Then

$$f(z) - f(z_0) = \operatorname{Re} 2\sqrt{s} \Big|_{z_0-1}^{z-1} + O(\varepsilon)$$

which yields (6.7).

Consider the case $z \in \mathbb{C}_-, z_0 \in \mathbb{C}_+$. The identity $f(z) = f(\bar{z})$ gives $f(z) - f(z_0) = f(\bar{z}) - f(z_0)$, $\bar{z}, z_0 \in \mathbb{C}_-$. Then the simple estimate $|\bar{z} - z_0| \leq |z - z_0|$ and the case $z, z_0 \in \overline{\mathbb{C}_+}$ imply (6.7).

Let $|z_0| > 2$. Let $a_0 = z_0/|z_0|, a = z/|z_0|$. Note that $|a - a_0| \leq \varepsilon$.

$$f(z) - f(z_0) = \operatorname{Re} \int_{z_0}^z \frac{dt}{\sqrt{t^2 - 1}} = \operatorname{Re} \int_{a_0}^a F(s) ds, \quad F(s) = \left(s^2 - \frac{1}{|z_0|^2} \right)^{-\frac{1}{2}}, \quad s = t/|z_0|.$$

We obtain

$$\left| s^2 - \frac{1}{|z_0|^2} \right| \geq |s|^2 - \frac{1}{4} \geq (1 - \varepsilon) - \frac{1}{4} \geq \frac{1}{2}, \quad \text{for } |s - a_0| \leq \varepsilon.$$

Thus the function $F(s) = \left(s^2 - \frac{1}{|z_0|^2} \right)^{-\frac{1}{2}}$ is analytic and bounded in $s, |a_0 - s| \leq \varepsilon$. Thus we deduce that $|f(z) - f(z_0)| \leq \varepsilon C$ for some absolute constant C . ■

Acknowledgments. The authors would like to thank Horst Hohberger for the Figure 2. Dmitry Chelkak was partly supported by grants VNP Minobrazovaniya 3.1–4733, RFFR 03–01–00377 and NSh–2266.2003.1. Evgeny Korotyaev was partly supported by DFG project BR691/23-1.

References

- [AS] Abramowitz, M. and Stegun, A., eds.: Handbook of Mathematical Functions. N.Y.: Dover Publications Inc., , 1992
- [Ah] Akhiezer, N.: The classical moment problem and some related questions in analysis. Hafner Publishing Co., New York 1965
- [BBK] Badanin, A; Brüning, J; Korotyaev, E. The Lyapunov function for Schrödinger operator with periodic 2×2 matrix potential, preprint 2005
- [BK] Badanin, A; Korotyaev, E. Spectral asymptotics for periodic forth order operators, will be published in Int. Math. Res. Not.
- [C1] R. Carlson. An inverse problem for the matrix Schrödinger equation. J. Math. Anal. Appl. 267 (2002), no. 2, 564–575.
- [C2] R. Carlson. Eigenvalue estimates and trace formulas for the matrix Hill's equation. J. Differential Equations 167 (2000), no. 1, 211–244.
- [C3] R. Carlson. Compactness of Floquet isospectral sets for the matrix Hill's equation. Proc. Amer. Math. Soc. 128 (2000), no. 10, 2933–2941.
- [C4] Carlson, R. A spectral transform for the matrix Hill's equation. Rocky Mountain J. Math. 34 (2004), no. 3, 869–895.
- [CG] Clark, S., Gesztesy, F.: Weyl-Titchmarsh M -function asymptotics, local uniqueness results, trace formulas, and Borg-type theorems for Dirac operators. Trans. Amer. Math. Soc. 354 (2002), no. 9, 3475–3534

- [CG1] S. Clark; F. Gesztesy. Weyl-Titchmarsh M -function asymptotics for matrix-valued Schrödinger operators. *Proc. London Math. Soc.* (3) 82 (2001), no. 3, 701–724.
- [CHGL] Clark S., Holden H., Gesztesy, F., Levitan, B.: Borg-type theorem for matrix-valued Schrödinger and Dirac operators, *J. Diff. Eqs.* 167(2000), 181–210
- [CS] Craig, W.; Simon, B. Subharmonicity of the Lyapunov index. *Duke Math. J.* 50 (1983), no. 2, 551–560.
- [DS] Dunford, N. and Schwartz, J.: *Linear Operators Part II: Spectral Theory*, Interscience, New York, 1988.
- [Fo] Forster O.: *Lectures on Riemann surfaces*. Graduate Texts in Mathematics, 81. Springer-Verlag, New York, 1991
- [GT1] J. Garnett, E. Trubowitz: Gaps and bands of one dimensional periodic Schrödinger operators. *Comment. Math. Helv.* 59, 258–312 (1984)
- [GT2] J. Garnett, E. Trubowitz: Gaps and bands of one dimensional periodic Schrödinger operators II. *Comment. Math. Helv.* 62, 18–37 (1987).
- [GL] Gel’fand I., Lidskii, V.: On the structure of the regions of stability of linear canonical systems of differential equations with periodic coefficients. (Russian) *Uspehi Mat. Nauk* (N.S.) 10 (1955), no. 1(63), 3–40.
- [Ge] Gel’fand, I.: Expansion in characteristic functions of an equation with periodic coefficients. (Russian) *Doklady Akad. Nauk SSSR* (N.S.) 73, (1950). 1117–1120.
- [GKM] Gesztesy, F., Kiselev A.; Makarov, K. Uniqueness results for matrix-valued Schrödinger, Jacobi, and Dirac-type operators. *Math. Nachr.* 239/240 (2002), 103–145.
- [Ka] T. Kappeler: Fibration of the phase space for the Korteweg-de-Vries equation. *Ann. Inst. Fourier (Grenoble)*, 41, 1, 539–575 (1991).
- [KK] Kargaev P., Korotyaev E.: Inverse Problem for the Hill Operator, the Direct Approach. *Invent. Math.*, 129(1997), no. 3, 567–593
- [KK1] Kargaev P., Korotyaev E.: Identities for the Dirichlet integral of subharmonic functions from the Cartright class, *Complex Var. Theory Appl.* 50 (2005), no. 1, 35–50.
- [KK2] Kargaev P., Korotyaev E.: Conformal mappings and subharmonic functions of Polya class
- [Ka] Kato, T.: *Perturbation theory for linear operators*. Springer-Verlag, Berlin, 1995.
- [K1] E. Korotyaev. The inverse problem for the Hill operator. I *Internat. Math. Res. Notices*, 3(1997), 113–125
- [K2] E. Korotyaev. Inverse problem and the trace formula for the Hill operator. II *Math. Z.* 231(1999), no. 2, 345–368
- [K3] E. Korotyaev. Inverse problem for periodic "weighted" operators, *J. Funct. Anal.* 170(2000), no. 1, 188–218
- [K4] E. Korotyaev. Marchenko-Ostrovski mapping for periodic Zakharov-Shabat systems, *J. Differential Equations*, 175(2001), no. 2, 244–274
- [K5] E. Korotyaev. Inverse Problem and Estimates for Periodic Zakharov-Shabat systems, *J. Reine Angew. Math.* 583(2005), 87–115
- [K6] E. Korotyaev. Characterization of the spectrum of Schrödinger operators with periodic distributions. *Int. Math. Res. Not.* (2003) no. 37, 2019–2031

- [K7] Korotyaev, E.: The estimates of periodic potentials in terms of effective masses. *Comm. Math. Phys.* 183 (1997), no. 2, 383–400.
- [K8] Korotyaev, E.: Metric properties of conformal mappings on the complex plane with parallel slits, *Internat. Math. Res. Notices*, 10(1996), 493–503
- [K9] Korotyaev, E.: Estimates for the Hill operator. I *J. Differential Equations* 162 (2000), no. 1, 1–26
- [K10] Korotyaev, E.: Estimates of periodic potentials in terms of gap lengths, *Comm. Math. Phys.* 197 (1998), no. 3, 521–526
- [K11] Korotyaev, E.: Estimates for the Hill operator, II, will be published in *J. Differential Equations*,
- [Kr] Krein, M.: The basic propositions of the theory of λ -zones of stability of a canonical system of linear differential equations with periodic coefficients. In memory of A. A. Andronov, pp. 413–498. *Izdat. Akad. Nauk SSSR*, Moscow, 1955.
- [Ly] Lyapunov, A.: The general problem of stability of motion, 2 nd ed. *Gl. Red. Obshchetekh. Lit.*, Leningrad, Moscow, 1935; reprint *Ann. Math. Studies*, no. 17, Princeton Univ. Press, Princeton, N.J. 1947
- [MV] Maksudov, F.; Veliev, O. Spectral analysis of differential operators with periodic matrix coefficients. (Russian) *Differentsial'nye Uravneniya* 25 (1989), no. 3, 400–409, 547; translation in *Differential Equations* 25 (1989), no. 3, 271–277
- [MO1] Marchenko V., Ostrovski I.: A characterization of the spectrum of the Hill operator. *Math. USSR Sb.* 26, 493–554 (1975).
- [MO2] Marchenko V., Ostrovski I.: Approximation of periodic by finite-zone potentials. *Selecta Math. Sovietica.* 1987, 6, No 2, 101–136.
- [Mi1] Misura T. Properties of the spectra of periodic and anti-periodic boundary value problems generated by Dirac operators. I,II, *Theor. Funktsii Funktsional. Anal. i Prilozhen*, (Russian), 30 (1978), 90–101; 31 (1979), 102–109
- [Mi2] Misura T. Finite-zone Dirac operators. *Theor. Funktsii Funktsional. Anal. i Prilozhen*, (Russian), 33 (1980), 107–11.
- [PT] Pöschel, J., Trubowitz, E.: *Inverse spectral theory.* Pure and Applied Mathematics, 130. Academic Press, Inc., Boston, MA, 1987. 192 pp.
- [RS] M. Reed ; B. Simon. *Methods of modern mathematical physics. IV. Analysis of operators.* Academic Press, New York-London, 1978
- [Sh] Shen, Chao-Liang: Some eigenvalue problems for the vectorial Hill's equation. *Inverse Problems* 16 (2000), no. 3, 749–783.
- [Sp] Springer, G.: *Introduction to Riemann surfaces.* Addison-Wesley Publishing Company, Inc., Reading, Mass. 1957
- [Ti] Titchmarsh, E.: *Eigenfunction expansions associated with second-order differential equations* 2, Oxford: Clarendon Press 1958
- [YS] Yakubovich, V., Starzhinskii, V.: *Linear differential equations with periodic coefficients.* 1, 2. Halsted Press [John Wiley & Sons] New York-Toronto, 1975. Vol. 1, Vol. 2